



## AP<sup>®</sup> Calculus BC 1998 Free-Response Questions

**These materials are intended for non-commercial use by AP teachers for course and exam preparation; permission for any other use must be sought from the Advanced Placement Program. Teachers may reproduce them, in whole or in part, in limited quantities, for face-to-face teaching purposes but may not mass distribute the materials, electronically or otherwise. These materials and any copies made of them may not be resold, and the copyright notices must be retained as they appear here. This permission does not apply to any third-party copyrights contained herein.**

These materials were produced by Educational Testing Service (ETS), which develops and administers the examinations of the Advanced Placement Program for the College Board. The College Board and Educational Testing Service (ETS) are dedicated to the principle of equal opportunity, and their programs, services, and employment policies are guided by that principle.

The College Board is a national nonprofit membership association dedicated to preparing, inspiring, and connecting students to college and opportunity. Founded in 1900, the association is composed of more than 4,200 schools, colleges, universities, and other educational organizations. Each year, the College Board serves over three million students and their parents, 22,000 high schools, and 3,500 colleges through major programs and services in college admission, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT<sup>®</sup>, the PSAT/NMSQT<sup>®</sup>, and the Advanced Placement Program<sup>®</sup> (AP<sup>®</sup>). The College Board is committed to the principles of equity and excellence, and that commitment is embodied in all of its programs, services, activities, and concerns.

For further information, contact [www.collegeboard.com](http://www.collegeboard.com).

Copyright ©2002 by College Entrance Examination Board. All rights reserved. College Board, Advanced Placement Program, AP, and the acorn logo are registered trademarks of the College Entrance Examination Board.

**1998 Calculus BC Free-Response Questions**

**CALCULUS BC**

**Section II**

Time—1 hour and 30 minutes

Number of problems—6

Percent of total grade—50

A GRAPHING CALCULATOR IS REQUIRED FOR SOME PROBLEMS OR PARTS OF PROBLEMS ON THIS SECTION OF THE EXAMINATION.

**REMEMBER TO SHOW YOUR SETUPS AS DESCRIBED IN THE GENERAL INSTRUCTIONS.**

- 
1. Let  $R$  be the region in the first quadrant bounded by the graph of  $y = 8 - x^{\frac{3}{2}}$ , the  $x$ -axis, and the  $y$ -axis.
- (a) Find the area of the region  $R$ .
  - (b) Find the volume of the solid generated when  $R$  is revolved about the  $x$ -axis.
  - (c) The vertical line  $x = k$  divides the region  $R$  into two regions such that when these two regions are revolved about the  $x$ -axis, they generate solids with equal volumes. Find the value of  $k$ .
- 

**GO ON TO THE NEXT PAGE** 

## 1998 Calculus BC Free-Response Questions

2. Let  $f$  be the function given by  $f(x) = 2xe^{2x}$ .
- (a) Find  $\lim_{x \rightarrow -\infty} f(x)$  and  $\lim_{x \rightarrow \infty} f(x)$ .
  - (b) Find the absolute minimum value of  $f$ . Justify that your answer is an absolute minimum.
  - (c) What is the range of  $f$ ?
  - (d) Consider the family of functions defined by  $y = bxe^{bx}$ , where  $b$  is a nonzero constant. Show that the absolute minimum value of  $bxe^{bx}$  is the same for all nonzero values of  $b$ .
- 

**GO ON TO THE NEXT PAGE** 

### 1998 Calculus BC Free-Response Questions


3. Let  $f$  be a function that has derivatives of all orders for all real numbers. Assume  $f(0) = 5$ ,  $f'(0) = -3$ ,  $f''(0) = 1$ , and  $f'''(0) = 4$ .
- (a) Write the third-degree Taylor polynomial for  $f$  about  $x = 0$  and use it to approximate  $f(0.2)$ .
  - (b) Write the fourth-degree Taylor polynomial for  $g$ , where  $g(x) = f(x^2)$ , about  $x = 0$ .
  - (c) Write the third-degree Taylor polynomial for  $h$ , where  $h(x) = \int_0^x f(t) dt$ , about  $x = 0$ .
  - (d) Let  $h$  be defined as in part (c). Given that  $f(1) = 3$ , either find the exact value of  $h(1)$  or explain why it cannot be determined.
- 

**GO ON TO THE NEXT PAGE** 

### 1998 Calculus BC Free-Response Questions

4. Consider the differential equation given by  $\frac{dy}{dx} = \frac{xy}{2}$ .

- (a) On the axes provided below, sketch a slope field for the given differential equation at the nine points indicated.



- (b) Let  $y = f(x)$  be the particular solution to the given differential equation with the initial condition  $f(0) = 3$ . Use Euler's method starting at  $x = 0$ , with a step size of 0.1, to approximate  $f(0.2)$ . Show the work that leads to your answer.
- (c) Find the particular solution  $y = f(x)$  to the given differential equation with the initial condition  $f(0) = 3$ . Use your solution to find  $f(0.2)$ .
- 

**GO ON TO THE NEXT PAGE** 


### 1998 Calculus BC Free-Response Questions

5. The temperature outside a house during a 24-hour period is given by

$$F(t) = 80 - 10 \cos\left(\frac{\pi t}{12}\right), \quad 0 \leq t \leq 24,$$

where  $F(t)$  is measured in degrees Fahrenheit and  $t$  is measured in hours.

- (a) Sketch the graph of  $F$  on the grid below.



- (b) Find the average temperature, to the nearest degree Fahrenheit, between  $t = 6$  and  $t = 14$ .
- (c) An air conditioner cooled the house whenever the outside temperature was at or above 78 degrees Fahrenheit. For what values of  $t$  was the air conditioner cooling the house?
- (d) The cost of cooling the house accumulates at the rate of \$0.05 per hour for each degree the outside temperature exceeds 78 degrees Fahrenheit. What was the total cost, to the nearest cent, to cool the house for this 24-hour period?
- 

**GO ON TO THE NEXT PAGE** 

**1998 Calculus BC Free-Response Questions**

6. A particle moves along the curve defined by the equation  $y = x^3 - 3x$ . The  $x$ -coordinate of the particle,  $x(t)$ , satisfies the equation  $\frac{dx}{dt} = \frac{1}{\sqrt{2t+1}}$ , for  $t \geq 0$  with initial condition  $x(0) = -4$ .
- (a) Find  $x(t)$  in terms of  $t$ .
  - (b) Find  $\frac{dy}{dt}$  in terms of  $t$ .
  - (c) Find the location and speed of the particle at time  $t = 4$ .
- 

END OF EXAMINATION