


AP[®] CALCULUS BC
2005 SCORING GUIDELINES

Question 1

Let f and g be the functions given by $f(x) = \frac{1}{4} + \sin(\pi x)$ and $g(x) = 4^{-x}$. Let

R be the shaded region in the first quadrant enclosed by the y -axis and the graphs of f and g , and let S be the shaded region in the first quadrant enclosed by the graphs of f and g , as shown in the figure above.



- (a) Find the area of R .
 (b) Find the area of S .
 (c) Find the volume of the solid generated when S is revolved about the horizontal line $y = -1$.

$$f(x) = g(x) \text{ when } \frac{1}{4} + \sin(\pi x) = 4^{-x}.$$

f and g intersect when $x = 0.178218$ and when $x = 1$.

Let $a = 0.178218$.

(a) $\int_0^a (g(x) - f(x)) dx = 0.064 \text{ or } 0.065$

3 : $\begin{cases} 1 : \text{limits} \\ 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

(b) $\int_a^1 (f(x) - g(x)) dx = 0.410$

3 : $\begin{cases} 1 : \text{limits} \\ 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$


(c) $\pi \int_a^1 ((f(x) + 1)^2 - (g(x) + 1)^2) dx = 4.558 \text{ or } 4.559$

3 : $\begin{cases} 2 : \text{integrand} \\ 1 : \text{limits, constant, and answer} \end{cases}$

**AP[®] CALCULUS BC
2005 SCORING GUIDELINES**

Question 2

The curve above is drawn in the xy -plane and is described by the equation in polar coordinates $r = \theta + \sin(2\theta)$ for $0 \leq \theta \leq \pi$, where r is measured in meters and θ is measured in radians. The derivative of r with respect to θ is given by $\frac{dr}{d\theta} = 1 + 2\cos(2\theta)$.



- (a) Find the area bounded by the curve and the x -axis.
- (b) Find the angle θ that corresponds to the point on the curve with x -coordinate -2 .
- (c) For $\frac{\pi}{3} < \theta < \frac{2\pi}{3}$, $\frac{dr}{d\theta}$ is negative. What does this fact say about r ? What does this fact say about the curve?
- (d) Find the value of θ in the interval $0 \leq \theta \leq \frac{\pi}{2}$ that corresponds to the point on the curve in the first quadrant with greatest distance from the origin. Justify your answer.

(a)
$$\text{Area} = \frac{1}{2} \int_0^\pi r^2 d\theta$$

$$= \frac{1}{2} \int_0^\pi (\theta + \sin(2\theta))^2 d\theta = 4.382$$

3 : $\begin{cases} 1 : \text{limits and constant} \\ 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

(b)
$$-2 = r \cos(\theta) = (\theta + \sin(2\theta)) \cos(\theta)$$

$$\theta = 2.786$$

2 : $\begin{cases} 1 : \text{equation} \\ 1 : \text{answer} \end{cases}$

(c) Since $\frac{dr}{d\theta} < 0$ for $\frac{\pi}{3} < \theta < \frac{2\pi}{3}$, r is decreasing on this interval. This means the curve is getting closer to the origin.

2 : $\begin{cases} 1 : \text{information about } r \\ 1 : \text{information about the curve} \end{cases}$

(d) The only value in $\left[0, \frac{\pi}{2}\right]$ where $\frac{dr}{d\theta} = 0$ is $\theta = \frac{\pi}{3}$.

2 : $\begin{cases} 1 : \theta = \frac{\pi}{3} \text{ or } 1.047 \\ 1 : \text{answer with justification} \end{cases}$

θ	r
0	0
$\frac{\pi}{3}$	1.913
$\frac{\pi}{2}$	1.571

The greatest distance occurs when $\theta = \frac{\pi}{3}$.

AP[®] CALCULUS BC
2005 SCORING GUIDELINES

Question 3

Distance x (cm)	0	1	5	6	8
Temperature $T(x)$ ($^{\circ}\text{C}$)	100	93	70	62	55

A metal wire of length 8 centimeters (cm) is heated at one end. The table above gives selected values of the temperature $T(x)$, in degrees Celsius ($^{\circ}\text{C}$), of the wire x cm from the heated end. The function T is decreasing and twice differentiable.

- (a) Estimate $T'(7)$. Show the work that leads to your answer. Indicate units of measure.
- (b) Write an integral expression in terms of $T(x)$ for the average temperature of the wire. Estimate the average temperature of the wire using a trapezoidal sum with the four subintervals indicated by the data in the table. Indicate units of measure.
- (c) Find $\int_0^8 T'(x) dx$, and indicate units of measure. Explain the meaning of $\int_0^8 T'(x) dx$ in terms of the temperature of the wire.
- (d) Are the data in the table consistent with the assertion that $T''(x) > 0$ for every x in the interval $0 < x < 8$? Explain your answer.

(a) $\frac{T(8) - T(6)}{8 - 6} = \frac{55 - 62}{2} = -\frac{7}{2}^{\circ}\text{C/cm}$

1 : answer

(b) $\frac{1}{8} \int_0^8 T(x) dx$

Trapezoidal approximation for $\int_0^8 T(x) dx$:

$$A = \frac{100 + 93}{2} \cdot 1 + \frac{93 + 70}{2} \cdot 4 + \frac{70 + 62}{2} \cdot 1 + \frac{62 + 55}{2} \cdot 2$$

Average temperature $\approx \frac{1}{8} A = 75.6875^{\circ}\text{C}$

3 : $\begin{cases} 1 : \frac{1}{8} \int_0^8 T(x) dx \\ 1 : \text{trapezoidal sum} \\ 1 : \text{answer} \end{cases}$

(c) $\int_0^8 T'(x) dx = T(8) - T(0) = 55 - 100 = -45^{\circ}\text{C}$

The temperature drops 45°C from the heated end of the wire to the other end of the wire.

2 : $\begin{cases} 1 : \text{value} \\ 1 : \text{meaning} \end{cases}$

(d) Average rate of change of temperature on $[1, 5]$ is $\frac{70 - 93}{5 - 1} = -5.75$.

Average rate of change of temperature on $[5, 6]$ is $\frac{62 - 70}{6 - 5} = -8$.

No. By the MVT, $T'(c_1) = -5.75$ for some c_1 in the interval $(1, 5)$ and $T'(c_2) = -8$ for some c_2 in the interval $(5, 6)$. It follows that T' must decrease somewhere in the interval (c_1, c_2) . Therefore T'' is not positive for every x in $[0, 8]$.

2 : $\begin{cases} 1 : \text{two slopes of secant lines} \\ 1 : \text{answer with explanation} \end{cases}$


Units of $^{\circ}\text{C/cm}$ in (a), and $^{\circ}\text{C}$ in (b) and (c)

1 : units in (a), (b), and (c)

**AP[®] CALCULUS BC
2005 SCORING GUIDELINES**


Question 4

Consider the differential equation $\frac{dy}{dx} = 2x - y$.



- (a) On the axes provided, sketch a slope field for the given differential equation at the twelve points indicated, and sketch the solution curve that passes through the point $(0, 1)$. (Note: Use the axes provided in the pink test booklet.)
- (b) The solution curve that passes through the point $(0, 1)$ has a local minimum at $x = \ln\left(\frac{3}{2}\right)$. What is the y -coordinate of this local minimum?
- (c) Let $y = f(x)$ be the particular solution to the given differential equation with the initial condition $f(0) = 1$. Use Euler's method, starting at $x = 0$ with two steps of equal size, to approximate $f(-0.4)$. Show the work that leads to your answer.
- (d) Find $\frac{d^2y}{dx^2}$ in terms of x and y . Determine whether the approximation found in part (c) is less than or greater than $f(-0.4)$. Explain your reasoning.

(a)



3 : $\begin{cases} 1 : \text{zero slopes} \\ 1 : \text{nonzero slopes} \\ 1 : \text{curve through } (0, 1) \end{cases}$

(b) $\frac{dy}{dx} = 0$ when $2x = y$

The y -coordinate is $2\ln\left(\frac{3}{2}\right)$.

2 : $\begin{cases} 1 : \text{sets } \frac{dy}{dx} = 0 \\ 1 : \text{answer} \end{cases}$

(c) $f(-0.2) \approx f(0) + f'(0)(-0.2)$
 $\quad \quad \quad = 1 + (-1)(-0.2) = 1.2$
 $f(-0.4) \approx f(-0.2) + f'(-0.2)(-0.2)$
 $\quad \quad \quad \approx 1.2 + (-1.6)(-0.2) = 1.52$

2 : $\begin{cases} 1 : \text{Euler's method with two steps} \\ 1 : \text{Euler approximation to } f(-0.4) \end{cases}$

(d) $\frac{d^2y}{dx^2} = 2 - \frac{dy}{dx} = 2 - 2x + y$

$\frac{d^2y}{dx^2}$ is positive in quadrant II because $x < 0$ and $y > 0$.


$1.52 < f(-0.4)$ since all solution curves in quadrant II are concave up.

2 : $\begin{cases} 1 : \frac{d^2y}{dx^2} \\ 1 : \text{answer with reason} \end{cases}$

AP[®] CALCULUS BC
2005 SCORING GUIDELINES

Question 5

A car is traveling on a straight road. For $0 \leq t \leq 24$ seconds, the car's velocity $v(t)$, in meters per second, is modeled by the piecewise-linear function defined by the graph above.



- (a) Find $\int_0^{24} v(t) \, dt$. Using correct units, explain the meaning of $\int_0^{24} v(t) \, dt$.
- (b) For each of $v'(4)$ and $v'(20)$, find the value or explain why it does not exist. Indicate units of measure.
- (c) Let $a(t)$ be the car's acceleration at time t , in meters per second per second. For $0 < t < 24$, write a piecewise-defined function for $a(t)$.
- (d) Find the average rate of change of v over the interval $8 \leq t \leq 20$. Does the Mean Value Theorem guarantee a value of c , for $8 < c < 20$, such that $v'(c)$ is equal to this average rate of change? Why or why not?

- (a) $\int_0^{24} v(t) \, dt = \frac{1}{2}(4)(20) + (12)(20) + \frac{1}{2}(8)(20) = 360$
 The car travels 360 meters in these 24 seconds.

2 : $\begin{cases} 1 : \text{value} \\ 1 : \text{meaning with units} \end{cases}$

- (b) $v'(4)$ does not exist because

$$\lim_{t \rightarrow 4^-} \left(\frac{v(t) - v(4)}{t - 4} \right) = 5 \neq 0 = \lim_{t \rightarrow 4^+} \left(\frac{v(t) - v(4)}{t - 4} \right).$$

$$v'(20) = \frac{20 - 0}{16 - 24} = -\frac{5}{2} \text{ m/sec}^2$$

3 : $\begin{cases} 1 : v'(4) \text{ does not exist, with explanation} \\ 1 : v'(20) \\ 1 : \text{units} \end{cases}$

- (c)
$$a(t) = \begin{cases} 5 & \text{if } 0 < t < 4 \\ 0 & \text{if } 4 < t < 16 \\ -\frac{5}{2} & \text{if } 16 < t < 24 \end{cases}$$

 $a(t)$ does not exist at $t = 4$ and $t = 16$.

2 : $\begin{cases} 1 : \text{finds the values } 5, 0, -\frac{5}{2} \\ 1 : \text{identifies constants with correct intervals} \end{cases}$

- (d) The average rate of change of v on $[8, 20]$ is

$$\frac{v(20) - v(8)}{20 - 8} = -\frac{5}{6} \text{ m/sec}^2.$$

 No, the Mean Value Theorem does not apply to v on $[8, 20]$ because v is not differentiable at $t = 16$.

2 : $\begin{cases} 1 : \text{average rate of change of } v \text{ on } [8, 20] \\ 1 : \text{answer with explanation} \end{cases}$

**AP[®] CALCULUS BC
2005 SCORING GUIDELINES**

Question 6

Let f be a function with derivatives of all orders and for which $f(2) = 7$. When n is odd, the n th derivative of f at $x = 2$ is 0. When n is even and $n \geq 2$, the n th derivative of f at $x = 2$ is given by $f^{(n)}(2) = \frac{(n-1)!}{3^n}$.

- (a) Write the sixth-degree Taylor polynomial for f about $x = 2$.
- (b) In the Taylor series for f about $x = 2$, what is the coefficient of $(x - 2)^{2n}$ for $n \geq 1$?
- (c) Find the interval of convergence of the Taylor series for f about $x = 2$. Show the work that leads to your answer.

(a) $P_6(x) = 7 + \frac{1!}{3^2} \cdot \frac{1}{2!}(x-2)^2 + \frac{3!}{3^4} \cdot \frac{1}{4!}(x-2)^4 + \frac{5!}{3^6} \cdot \frac{1}{6!}(x-2)^6$

3 : $\left\{ \begin{array}{l} 1 : \text{polynomial about } x = 2 \\ 2 : P_6(x) \\ \langle -1 \rangle \text{ each incorrect term} \\ \langle -1 \rangle \text{ max for all extra terms,} \\ \quad + \dots, \text{ misuse of equality} \end{array} \right.$

(b) $\frac{(2n-1)!}{3^{2n}} \cdot \frac{1}{(2n)!} = \frac{1}{3^{2n}(2n)}$

1 : coefficient

- (c) The Taylor series for f about $x = 2$ is

$$f(x) = 7 + \sum_{n=1}^{\infty} \frac{1}{2n \cdot 3^{2n}} (x-2)^{2n}.$$

$$L = \lim_{n \rightarrow \infty} \left| \frac{\frac{1}{2(n+1)} \cdot \frac{1}{3^{2(n+1)}} (x-2)^{2(n+1)}}{\frac{1}{2n} \cdot \frac{1}{3^{2n}} (x-2)^{2n}} \right|$$

$$= \lim_{n \rightarrow \infty} \left| \frac{2n}{2(n+1)} \cdot \frac{3^{2n}}{3^2 3^{2n}} (x-2)^2 \right| = \frac{(x-2)^2}{9}$$

$$L < 1 \text{ when } |x-2| < 3.$$

Thus, the series converges when $-1 < x < 5$.

When $x = 5$, the series is $7 + \sum_{n=1}^{\infty} \frac{3^{2n}}{2n \cdot 3^{2n}} = 7 + \frac{1}{2} \sum_{n=1}^{\infty} \frac{1}{n},$

which diverges, because $\sum_{n=1}^{\infty} \frac{1}{n}$, the harmonic series, diverges.

When $x = -1$, the series is $7 + \sum_{n=1}^{\infty} \frac{(-3)^{2n}}{2n \cdot 3^{2n}} = 7 + \frac{1}{2} \sum_{n=1}^{\infty} \frac{1}{n},$

which diverges, because $\sum_{n=1}^{\infty} \frac{1}{n}$, the harmonic series, diverges.

The interval of convergence is $(-1, 5)$.

5 : $\left\{ \begin{array}{l} 1 : \text{sets up ratio} \\ 1 : \text{computes limit of ratio} \\ 1 : \text{identifies interior of} \\ \quad \text{interval of convergence} \\ 1 : \text{considers both endpoints} \\ 1 : \text{analysis/conclusion for} \\ \quad \text{both endpoints} \end{array} \right.$