



## **AP<sup>®</sup> Calculus BC 2005 Sample Student Responses**

### **The College Board: Connecting Students to College Success**

The College Board is a not-for-profit membership association whose mission is to connect students to college success and opportunity. Founded in 1900, the association is composed of more than 4,700 schools, colleges, universities, and other educational organizations. Each year, the College Board serves over three and a half million students and their parents, 23,000 high schools, and 3,500 colleges through major programs and services in college admissions, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT<sup>®</sup>, the PSAT/NMSQT<sup>®</sup>, and the Advanced Placement Program<sup>®</sup> (AP<sup>®</sup>). The College Board is committed to the principles of excellence and equity, and that commitment is embodied in all of its programs, services, activities, and concerns.


Copyright © 2005 by College Board. All rights reserved. College Board, AP Central, APCD, Advanced Placement Program, AP, AP Vertical Teams, Pre-AP, SAT, and the acorn logo are registered trademarks of the College Entrance Examination Board. Admitted Class Evaluation Service, CollegeEd, Connect to college success, MyRoad, SAT Professional Development, SAT Readiness Program, and Setting the Cornerstones are trademarks owned by the College Entrance Examination Board. PSAT/NMSQT is a registered trademark of the College Entrance Examination Board and National Merit Scholarship Corporation. Other products and services may be trademarks of their respective owners. Permission to use copyrighted College Board materials may be requested online at: <http://www.collegeboard.com/inquiry/cbpermit.html>.

**Visit the College Board on the Web: [www.collegeboard.com](http://www.collegeboard.com).**

**AP Central is the official online home for the AP Program and Pre-AP: [apcentral.collegeboard.com](http://apcentral.collegeboard.com).**

CALCULUS AB  
SECTION II, Part A  
Time—45 minutes  
Number of problems—3

A graphing calculator is required for some problems or parts of problems.



Work for problem 1(a)

$$f(x) = \frac{1}{4} + \sin(\pi x)$$

$$g(x) = 4^{-x}$$

$$A_R = \int_0^{-.178} g(x) - f(x) \, dx$$

$$A_R = \int_0^{-.178} 4^{-x} - \frac{1}{4} - \sin(\pi x) \, dx$$

$$A_R = .0648 \text{ u}^2$$

Do not write beyond this border.

Continue problem 1 on page 5.


Work for problem 1(b)

$$A_S = \int_{-.178}^1 f(x) - g(x) dx$$

$$A_S = \int_{-.178}^1 \frac{1}{4} + \sin(\pi x) - 4^{-x} dx$$

$$A_S = .410 u^2$$

Work for problem 1(c)



$$R = f(x) - -1$$

$$r = g(x) - -1$$

$$V_S = \int_{-.178}^1 \pi R^2 - \pi r^2 dx$$

$$V_S = \int_{-.178}^1 \pi \left( \frac{1}{4} + \sin(\pi x) + 1 \right)^2 - \pi \left( 4^{-x} + 1 \right)^2 dx$$

$$V_S = 4.559 u^3$$

Do not write beyond this border.


GO ON TO THE NEXT PAGE.

CALCULUS AB  
SECTION II, Part A

Time—45 minutes

Number of problems—3

A graphing calculator is required for some problems or parts of problems.



Work for problem 1(a)

$$\int_{0.17821865}^{1.94167} g(x) - f(x) dx$$

$$= 0.065$$

Do not write beyond this border.

Continue problem 1 on page 5.

Work for problem 1(b)

$$\int_{-1.7821865}^{1.94167} f(x) - g(x) dx$$
$$= 0.117$$

Work for problem 1(c)

$$\int_{-1.7821865}^{1.94167} (-1 - f(x))^2 - (-1 - g(x))^2 dx$$
$$= 0.618$$

Do not write beyond this border.


GO ON TO THE NEXT PAGE.

CALCULUS AB  
SECTION II, Part A

Time—45 minutes

Number of problems—3

A graphing calculator is required for some problems or parts of problems.



Work for problem 1(a)

$$\int_0^{0.781} 4^{-x} - \left[ \frac{1}{4} + \sin(\pi x) \right] dx$$

Area of R = -0.2824 units<sup>2</sup>

intersect  
0.781

↑  
via TI-83

Do not write beyond this border.


Continue problem 1 on page 5.

Work for problem 1(b)

$$\int_{0.781}^1 f(x) - g(x) = \int_{0.781}^1 \left[ \frac{1}{4} + \sin(\pi x) \right] - 4^{-x}$$

Area of  $S = 0.0632 \text{ units}^2$  ← via TI-83

Work for problem 1(c)




$$\pi \int_{0.781}^1 \left[ (f(x))^2 - (g(x))^2 \right] dx$$

$$\pi \int_{0.781}^1 \left[ \left( \frac{1}{4} + \sin(\pi x) \right)^2 - 4^{-2x} \right] dx$$

Volume = -14.5243 units<sup>3</sup>

Do not write beyond this border.

GO ON TO THE NEXT PAGE.





Work for problem 2(a)

Polar Area =  $\frac{1}{2} \int r^2 d\theta$        $r^2 = (\theta + \sin(2\theta))^2$

$$\frac{1}{2} \int_0^{\pi} (\theta + \sin(2\theta))^2 d\theta = \underline{4.382 \text{ units}^2}$$

Do not write beyond this border.

Work for problem 2(b)



$$\theta + \sin 2\theta = r = \frac{x}{\cos \theta} \quad | \quad x = -2$$

$$r \cos \theta = x$$

$$r = \frac{x}{\cos \theta}$$

$$\theta + \sin 2\theta = \frac{-2}{\cos \theta}$$

$$\theta = \underline{2.786 \text{ radians}}$$

~~2.786~~      2.786

Continue problem 2 on page 7.

Work for problem 2(c)

This means that the length of  $r$  is decreasing w/ respect to  $\theta$ . Therefore the distance of the graph from the origin is getting smaller @ from  $\frac{\pi}{3}$  @  $\frac{2\pi}{3}$ .

Work for problem 2(d)

trying to maximize  $r$

$$r(\theta) = \theta + \sin(2\theta)$$

$$r'(\theta) = 1 + 2\cos(2\theta)$$


$$r'(\theta) = 0 \text{ when } \theta = \pi/3$$

$$\cos(2\theta) = -1/2$$

$$\frac{\pi}{3}$$

$$2\theta = 2\pi/3$$

$$\theta = \frac{2\pi}{6} = \frac{\pi}{3}$$




since  $r$  is increasing from interval is positive  $0 < \theta < \pi/3$  since  $r'(\theta) > 0$  that

since  $r$  is decreasing after  $\pi/3 > \theta > \pi/3$  since  $r'(\theta) < 0$  that interval is negative

when  $\theta = \pi/3$ ,  $r$  is max  $\therefore$  there is a max at  $\theta = \pi/3$

GO ON TO THE NEXT PAGE.

Do not write beyond this border.



Work for problem 2(a)

The shown area is from  $0 \leq \theta \leq \pi$ , so the area is given by:

$$\text{area} = \int_0^{\pi} \frac{1}{2} (\theta + \sin(2\theta))^2 d\theta = \boxed{4.382}$$

Work for problem 2(b)

$$x = r \cos \theta$$

$$= (\theta + \sin(2\theta))(\cos \theta)$$

Therefore, if  $x = -2$ ,

$$-2 = (\theta + \sin(2\theta))(\cos \theta)$$

$$\therefore \theta = \boxed{2.786}$$

Do not write beyond this border.

Do not write beyond this border.

Continue problem 2 on page 7.

2

2

2

2

2

2

2

2

2

2

B<sub>2</sub>

Work for problem 2(c)

Since from  $\frac{\pi}{3} < \theta < \frac{2\pi}{3}$   $\frac{dr}{d\theta} < 0$ ,  $r$  is steadily decreasing in that range. Since this range is between 0 and  $\pi$ , a range in which a constant- $r$  function will be concave down, and since this decreases even more than that does because  $r$  is itself diminishing, the curve is concave down in that range.


Work for problem 2(d)

If a point has the greatest distance from the origin,  $r$  must be at a maximum. For  $r = \theta + \sin(2\theta)$  from  $0 \leq \theta \leq \frac{\pi}{2}$ ,  $r$  reaches a maximum when  $\theta$  is 1.047. At that angle, the curve is thus its furthest from the origin in the first quadrant.

Do not write beyond this border.

Do not write beyond this border.

GO ON TO THE NEXT PAGE.



Work for problem 2(a)

$$A = \frac{1}{2} \int_0^{\pi} (\theta + \sin(2\theta))^2 d\theta$$

$$= 8.765$$

Work for problem 2(b)

$$x = r \cos \theta$$

$$-z = r \cos \theta$$

Continue problem 2 on page 7.

Work for problem 2(c)

This says that as  $\theta$  increases during the interval from  $\frac{\pi}{3}$  to  $\frac{2\pi}{3}$  the length of  $r$  is decreasing.

It says the curve is bending towards itself more. Because  $r$  is decreasing, the curve is forming a tighter loop which also decreases area.


Work for problem 2(d)

$$r = \theta + \sin(2\theta)$$

$$\frac{dr}{d\theta} = 1 + 2\cos(2\theta)$$

$$0 = 1 + 2\cos(2\theta)$$

$$\theta = 1.1047$$



A maximum occurs at  $\theta = 1.1047$  because  $\frac{dr}{d\theta}$  changes from positive to negative.

A max. value of  $R$  occurs at  $\theta = 1.1047$

Do not write beyond this border.

Do not write beyond this border.

Distance $x$ (cm)	0	1	5	6	8
Temperature $T(x)$ ( $^{\circ}\text{C}$ )	100	93	70	62	55

Work for problem 3(a)

$$T'(x) = \frac{55 - 62}{8 - 6} = -3.5 \text{ } ^{\circ}\text{C}/\text{cm}$$

Work for problem 3(b)

$$\text{Average Temp} = \frac{1}{8} \int_0^8 T(x) dx.$$

$$\begin{aligned} \text{Average} &= \frac{1}{8} \cdot \left[ (100 + 93)(1)\left(\frac{1}{2}\right) + (93 + 70)(4)\left(\frac{1}{2}\right) \right. \\ &\quad \left. + (62 + 70)(1)\left(\frac{1}{2}\right) + (55 + 62)(2)\left(\frac{1}{2}\right) \right] \\ &= 75.688 \text{ } ^{\circ}\text{C} \end{aligned}$$

Do not write beyond this border.

Continue problem 3 on page 9.

Work for problem 3(c)

$$\begin{aligned}\int_0^8 T'(x) dx &= T(8) - T(0) \\ &= 55 - 100 \\ &= -45^\circ\text{C}\end{aligned}$$

$\int_0^8 T'(x) dx$  mean the total change (drop) in temperature of the wire from 0 cm to 8 cm.

Work for problem 3(d)

$T''(x) > 0 \Rightarrow T'(x)$  is increasing over the period.

from  $x = 0$  to 1

slope  $\Rightarrow -7$

$x = 1$  to 5

slope  $\Rightarrow \frac{70-93}{5-1} = -5.75$

$x = 5$  to 6

slope  $\Rightarrow \frac{62-70}{6-5} = -8$

$x = 6$  to 8

slope  $\Rightarrow \frac{55-62}{8-6} = -3.5$

By MVT.

$\therefore$  between 5 to 6 there is a point with slope  $-8$  which means a decrease of  $T'(x)$

$\Downarrow$   
 $T'(x)$  is not always increasing

$\therefore T''(x) > 0$  is not consistent in

END OF PART A OF SECTION II the table data.

IF YOU FINISH BEFORE TIME IS CALLED, YOU MAY CHECK YOUR WORK ON PART A ONLY. DO NOT GO ON TO PART B UNTIL YOU ARE TOLD TO DO SO.

Do not write beyond this border.

Distance $x$ (cm)	0	1	5	6	8
Temperature $T(x)$ ( $^{\circ}\text{C}$ )	100	93	70	62	55

Work for problem 3(a)

$$\begin{aligned}
 T'(7) &\approx \frac{T(8) - T(6)}{8 - 6} \\
 &\approx \frac{55 - 62}{2} \\
 &\approx -\frac{7}{2} \text{ } ^{\circ}\text{C/cm}
 \end{aligned}$$

Work for problem 3(b)

$$\begin{aligned}
 \text{Avg.} &= \frac{1}{8} \int_0^8 T(x) dx \\
 &\approx \frac{1}{8} \left[ \frac{1}{2}(T(0) + T(1)) + \frac{1}{2}(T(1) + T(5)) + \frac{1}{2}(T(5) + T(6)) + \frac{1}{2}(T(6) + T(8)) \right] \\
 &\approx \frac{1}{8} \left[ \frac{1}{2}(100 + 93) + \frac{1}{2}(93 + 70) + \frac{1}{2}(70 + 62) + \frac{1}{2}(62 + 55) \right] \\
 &\approx 37.813 \text{ } ^{\circ}\text{C}
 \end{aligned}$$

Do not write beyond this border.

Continue problem 3 on page 9.

Work for problem 3(c)

$$\begin{aligned}\int_0^8 T'(x) dx &= T(8) - T(0) \\ &= 55 - 100 \\ &= -45^\circ\text{C}\end{aligned}$$

This is the total change in temperature of the wire, from one end to the other.

Work for problem 3(d)

$$\begin{aligned}T'(0.5) &\cong \frac{T(1) - T(0)}{1 - 0} \\ &\cong -7.00\end{aligned}$$

$$\begin{aligned}T'(3) &\cong \frac{T(5) - T(1)}{5 - 1} \\ &\cong -5.750\end{aligned}$$

$$T'(7) \cong -3.500$$

The table is consistent with the assertion that  $T''(x) > 0$  for every  $x$  in the interval  $0 < x < 8$ , since  $T'(x)$  is increasing.

END OF PART A OF SECTION II

IF YOU FINISH BEFORE TIME IS CALLED, YOU MAY CHECK YOUR WORK ON PART A ONLY. DO NOT GO ON TO PART B UNTIL YOU ARE TOLD TO DO SO.

Do not write beyond this border.

Distance $x$ (cm)	0	1	5	6	8
Temperature $T(x)$ ( $^{\circ}\text{C}$ )	100	93	70	62	55

Work for problem 3(a)

$$\begin{aligned}
 T'(7) &\approx \frac{T(8) - T(6)}{8 - 6} \\
 &\approx \frac{55 - 62}{2} \\
 &\approx -7/2 \text{ } ^{\circ}\text{C/cm}
 \end{aligned}$$

Work for problem 3(b)

$$\text{Avg } T = \frac{\int_0^8 T(x) dx}{8}$$

$$\int_0^8 T(x) dx \approx \frac{b-a}{2n} [f(0) + 2f(1) + 2f(5) + 2f(6) + f(8)]$$

$$\approx \frac{8}{2(4)} [100 + 2(93) + 2(70) + 2(62) + 55]$$

$$\approx 605$$

$$\frac{\int_0^8 T(x) dx}{8} \approx \frac{605}{8} = \boxed{75.625 \text{ } ^{\circ}\text{C}}$$

Do not write beyond this border.

Continue problem 3 on page 9.

Work for problem 3(c)

$$\begin{aligned}\int_0^8 T'(x) dx &= T(x) \Big|_0^8 \\ &= T(8) - T(0) \\ &= 55 - 100 \\ &= -45^\circ\text{C/cm}\end{aligned}$$

$\int_0^8 T'(x) dx$  represents the average rate of change of the temperature of the wire as  $x$  increases from 0 to 8.

Work for problem 3(d)

$T''(x) > 0$  implies that  $T(x)$  is concave up, or that the rate of change is increasing.

The data in the table do not show that the rate of change is increasing from  $x=0$  to  $x=8$ . For example: from  $x=0$  to  $x=1$ ,  $T(x)$  decreases  $7^\circ\text{C}$ . In order for  $T(x)$  to be concave up, it must decrease by less than  $7^\circ\text{C/cm}$  from  $x=1$  to  $x=5$ :  $\frac{T(5) - T(1)}{4} = -5.75^\circ\text{C}$ .

$T(x)$  decreases by  $5.75^\circ\text{C}$ , which is less than  $7^\circ\text{C}$ , so it is changing at an increasing rate.

Therefore  $T(x)$  is concave up and  $T''(x) > 0$  is true for  $x=0$  to  $x=5$ .

$\frac{T(6) - T(5)}{1} = -10$  ← This is not consistent however, so  $T(x)$  is not concave up for all  $x$  in  $0 < x < 8$ .

END OF PART A OF SECTION II

IF YOU FINISH BEFORE TIME IS CALLED, YOU MAY CHECK YOUR WORK ON PART A ONLY. DO NOT GO ON TO PART B UNTIL YOU ARE TOLD TO DO SO.

Do not write beyond this border.

4 4 4 4 4 4 4 4 4 4

NO CALCULATOR ALLOWED

CALCULUS BC


SECTION II, Part B

Time—45 minutes

Number of problems—3

No calculator is allowed for these problems.

Work for problem 4(a)



Work for problem 4(b)

Local extrema:  $\frac{dy}{dx} = 0$ 

$$0 = 2 \ln\left(\frac{x}{2}\right) - y$$

$$y = 2 \ln\left(\frac{x}{2}\right)$$

Do not write beyond this border.

Continue problem 4 on page 11.

4 4 4 4 4 4 4 4 4 4

NO CALCULATOR ALLOWED

Work for problem 4(c)

$(x, y)$	$\Delta x$	$\Delta y = \Delta x \cdot (2x - y)$	next $(x, y)$
$(0, 1)$	$-0.2$	$-0.2(-1) = 0.2$	$(-0.2, 1.2)$
$(-0.2, 1.2)$	$-0.2$	$-0.2(-0.4 - 1.2) = 0.32$	$(-0.4, 1.52)$

↓

$f(-0.4) = \boxed{1.52}$

Do not write beyond this border.

Work for problem 4(d)

$$\frac{d^2 y}{dx^2} = 2 - \frac{dy}{dx} = 2 - 2x + y$$

$x$  negative &  $y$  positive for values in question;

$\frac{d^2 y}{dx^2}$  positive, graph concave up



The estimation is less than the actual value. The slopes used were greater than the actual slopes; when working from right to left, this means a lesser value.

GO ON TO THE NEXT PAGE.

4

4

4

4

4

4

4

4

4

4

NO CALCULATOR ALLOWED

CALCULUS BC


SECTION II, Part B

Time—45 minutes

Number of problems—3

No calculator is allowed for these problems.

Work for problem 4(a)



Work for problem 4(b)

$$\frac{dy}{dx} = 2\left(\ln\left(\frac{3}{2}\right)\right) - y$$

$$2\left(\ln\left(\frac{3}{2}\right)\right) - y = 0$$

$$-y = -2\left(\ln\left(\frac{3}{2}\right)\right)$$

$$y = 2\left(\ln\left(\frac{3}{2}\right)\right)$$

Continue problem 4 on page 11.



NO CALCULATOR ALLOWED

Work for problem 4(c)

$$\Delta x = -0.2$$

$$(0, 1) \quad \frac{dy}{dx} = 2x - y$$

$$= 2(0) - 1 = -1$$

$$\Delta y = -1(-0.2) = 0.2$$

$$(-0.2, 1.2)$$

$$2(-0.2) - 1.2 = -0.4 - 1.2 = -1.6$$

$$\Delta y = -1.6(-0.2) = 0.32$$

$$(-0.4, 1.04)$$

$$f(-0.4) = 1.04$$

Work for problem 4(d)

$$\frac{d^2y}{dx^2} = 2 - \frac{dy}{dx}$$

$$\frac{d^2y}{dx^2} = 2 - (2x - y)$$

$$\frac{d^2y}{dx^2} = 2 - 2x - y$$

Do not write beyond this border.

GO ON TO THE NEXT PAGE.

4

4

4

4

4

4

4

4

4

4

NO CALCULATOR ALLOWED

CALCULUS BC


SECTION II, Part B

Time—45 minutes

Number of problems—3

No calculator is allowed for these problems.

Work for problem 4(a)



Work for problem 4(b)

$$\frac{dy}{dx} = 2x - 5$$

$$\int \frac{1}{y} dy = \int 2x dx$$

$$-\ln y = x^2 + C$$

$$-\ln 1 = 0 + C$$

$$C = -\ln 1$$

$$-\ln 5 = e^2 - \ln 1$$

$$-\ln 5 = \left(\ln \frac{3}{2}\right)^2 - \ln 1$$

$$-\ln 5 = \ln \frac{9}{4}$$

$$\frac{9}{4} = 5$$

Do not write beyond this border.

Do not write beyond this border.

Continue problem 4 on page 11.

NO CALCULATOR ALLOWED

Work for problem 4(c)

$x$	$y$	$dx$	$dy$
0	1	-0.2	-0.2
-0.2	1.2	-0.2	-0.32
-0.4	1.52		

$$2x - y = \frac{dy}{dx}$$

$$2(0) - 1 = \frac{dy}{dx}$$

$$2(-0.2) - 1.2 = \frac{dy}{dx}$$

$$dx - 2x - y = dy$$

1.52


Do not write beyond this border.

Do not write beyond this border.

Work for problem 4(d)

$$\frac{d^2y}{dx^2} = 2 - y'$$

$$y' = 2x - y$$



$$\frac{d^2y}{dx^2} = 2 - 2x - y$$

less because the graph is concave up for  $(-0.4, 0)$  that means when it is estimated from the tangent line at  $0$ , it'll be less than actual

GO ON TO THE NEXT PAGE.

5

5

5

5

5

5

5


5

5

5

A

NO CALCULATOR ALLOWED



Work for problem 5(a)

$$\int_0^{24} v(t) dt = \frac{1}{2} (12 + 24)(20) = \boxed{360 \text{ meters}}$$

$\int_0^{24} v(t) dt$  is the displacement of the car in meters from time  $t=0$  seconds to  $t=24$  seconds - since the integral is positive, the car is 360 meters in the positive direction at time 24 seconds as compared with its position at time 0 seconds

Work for problem 5(b)

$v'(4)$  does not exist because  $\lim_{x \rightarrow 4^-} v'(x) = 5 \neq 0 = \lim_{x \rightarrow 4^+} v'(x)$

$$v'(20) = \frac{-20}{8} = \boxed{\frac{-5}{2} \frac{\text{meters}}{\text{second}^2}}$$

Do not write beyond this border.

Continue problem 5 on page 13.

NO CALCULATOR ALLOWED

Work for problem 5(c)

$$a(t) = \begin{cases} 5 \text{ m/s}^2 & \text{for } 0 \leq t < 4 \text{ seconds} \\ 0 \text{ m/s}^2 & \text{for } 4 < t < 16 \text{ seconds} \\ -5/2 \text{ m/s}^2 & \text{for } 16 < t \leq 24 \text{ seconds} \end{cases}$$

\*  $a(t)$  undefined for  $t = 4, 16$  seconds

Work for problem 5(d)


$$\frac{10 - 20}{20 - 8} = \frac{-10}{12} = \boxed{\frac{-5}{6} \frac{\text{m}}{\text{s}^2}}$$

The Mean Value Theorem does NOT guarantee a value for  $c$ ,  $8 < c < 20$ , such that  $v'(c)$  equals this average rate of change because  $v(t)$  does not fulfill the requirements for the Mean Value Theorem since  $v(t)$  is not differentiable on the interval  $(8, 20)$  since  $v'(t)$  is undefined at  $t = 16$  seconds.

Do not write beyond this border.

GO ON TO THE NEXT PAGE.

## NO CALCULATOR ALLOWED



Work for problem 5(a)

$$\frac{1}{2}(4)(20) + 12(20) + \frac{1}{2}(8)(20)$$

$$40 + 240 + 80 = 360 = \int_0^{24} v(t) dt$$

$\int_0^{24} v(t) dt = 360$  meters which is the total distance the car traveled from  $t=0$  to  $t=24$ .

Work for problem 5(b)

The derivative of  $v$  at  $t=4$  does not exist because it is located at a corner.

$$v'(20) = \frac{20-0}{16-24} = \frac{20}{-8} = -\frac{5}{2} \frac{\text{meters}}{\text{sec}}$$

Do not write beyond this border.

Continue problem 5 on page 13.

## NO CALCULATOR ALLOWED

Work for problem 5(c)

slope of  $v(t)$  from  $t=0$  to  $t=4$  : 5  
 slope " "  $t=4$  to  $t=16$  : 0  
 slope " "  $t=16$  to  $t=24$  :  $-\frac{5}{2}$

$$a(t) = \begin{cases} 5, & 0 \leq t \leq 4 \\ 0, & 4 < t \leq 16 \\ -\frac{5}{2}, & 16 < t \leq 24 \end{cases}$$

Work for problem 5(d)

The Mean Value Theorem does not guarantee this because  $v(t)$  is not differentiable over  $8 \leq t \leq 20$ .

~~$$\frac{\int_8^{20} (v(20) - v(8)) dt}{20-8} = \frac{(10-20) dt}{12} = \frac{\int_8^{20} -10 dt}{12}$$~~

$$4(20) + 4(20) + \frac{1}{2}(4)(20+10) = 80 + 80 + 60 = 220$$


$$\frac{220}{12} \frac{\text{meters}}{\text{sec}^2}$$

$$\frac{160}{50} = 220$$

Do not write beyond this border.

GO ON TO THE NEXT PAGE.


## NO CALCULATOR ALLOWED



Work for problem 5(a)

$\int_0^{24} (v(t)) dt$  is asking for the total distance traveled in these 24 seconds. This is the total Area underneath the graph.

$$\begin{aligned}
 A &= \frac{1}{2} (4)(20) + 12(20) + \frac{1}{2} (8)(20) \\
 &= 40 + 240 + 80 \\
 &= 360
 \end{aligned}$$



Work for problem 5(b)

$v'(t)$  is asking for the slope at a certain point in time. At  $t=4$  and  $t=20$  the slope is undefined because there is a corner at both of these times.

Do not write beyond this border.

Continue problem 5 on page 13.

NO CALCULATOR ALLOWED

Work for problem 5(c)

$$A(t) = v'(t)$$

$$v(t) = \begin{cases} 5x & 0 \leq x \leq 4 \\ 20 & 4 < x \leq 16 \\ -\frac{5}{2}x + 60 & 16 < x \leq 24 \end{cases}$$

$$A(t) = \begin{cases} 5 & 0 \leq x \leq 4 \\ 0 & 4 < x \leq 16 \\ -\frac{5}{2} & 16 < x \leq 24 \end{cases}$$

$\leftarrow 5$  is the deriv of  $5x$   
 $\leftarrow 0$  is the deriv of  $20$   
 $\leftarrow -\frac{5}{2}$  is the deriv of  $-\frac{5}{2}x + 60$

Do not write beyond this border.

Work for problem 5(d)

$$\text{avg rate of chg} = \frac{f(b) - f(a)}{b - a} = \frac{f(20) - f(8)}{20 - 8} = \frac{10 - 20}{20 - 8} = \boxed{-\frac{5}{6}}$$

• yes C has a guaranteed value because  $v(t)$  is continuous from  $8 \leq x \leq 20$ .

GO ON TO THE NEXT PAGE.

NO CALCULATOR ALLOWED

Work for problem 6(a)

$$\begin{array}{r} 81 \\ \times 3 \\ \hline 243 \end{array}$$

$$f(2) = 7$$

$$f'(2) = f''(2) = f'''(2) = 0$$

$$f^{(4)}(2) = \frac{(2-1)!}{3^2} = \frac{1}{9}$$


$$f^{(5)}(2) = \frac{(4-1)!}{3^4} = \frac{6}{81} = \frac{2}{27}$$

$$f^{(6)}(2) = \frac{(6-1)!}{3^6} = \frac{5 \cdot 4 \cdot 3 \cdot 2}{3^6} = \frac{40}{243}$$

$$P_6(x) = 7 + \frac{1/9(x-2)^2}{2!} + \frac{2/27(x-2)^4}{4!} + \frac{40/243(x-2)^6}{6!}$$

Work for problem 6(b)

$$\text{coefficient} = \frac{(2n-1)!}{3^{2n}} \cdot \frac{1}{(2n)!} = \frac{(2n-1)!}{(3^{2n})(2n)!} = \frac{1}{2n(3^{2n})}$$



Do not write beyond this border.

Continue problem 6 on page 15.

## NO CALCULATOR ALLOWED

Work for problem 6(c)

$$1 + \sum_{n=1}^{\infty} \frac{(x-2)^{2n}}{2n(9^n)}$$

$$\lim_{n \rightarrow \infty} \left| \frac{(x-2)^{2n+2}}{(2n+2)(9^{n+1})} \cdot \frac{2n(9^n)}{(x-2)^{2n}} \right|$$

$$\lim_{n \rightarrow \infty} \left| \frac{2n}{2n+2} \cdot \frac{(x-2)^2}{9} \right|$$

$$\left| \frac{(x-2)^2}{9} \right| < 1$$

$$(x-2)^2 < 9$$

$$-3 < x-2 < 3$$

$$-1 < x < 5$$

$$\text{If } x = -1: \sum_{n=1}^{\infty} \frac{(-3)^{2n}}{2n(3^{2n})} = \sum_{n=1}^{\infty} \frac{(-1)^{2n}}{2n} = \sum_{n=1}^{\infty} \frac{1}{2n}$$

$\frac{1}{n}$  = harmonic series  
 $\therefore$  DN

$$\text{If } x = 5: \sum_{n=1}^{\infty} \frac{3^{2n}}{2n(3^{2n})} = \sum_{n=1}^{\infty} \frac{1}{2} \cdot \frac{1}{n}$$

harmonic series  
 $\therefore$  DIV

END OF EXAM

THE FOLLOWING INSTRUCTIONS APPLY TO THE BACK COVER OF THIS SECTION II BOOKLET.

- MAKE SURE YOU HAVE COMPLETED THE IDENTIFICATION INFORMATION AS REQUESTED ON THE BACK OF THIS SECTION II BOOKLET.
- CHECK TO SEE THAT YOUR AP NUMBER APPEARS IN THE BOX(ES) ON THE BACK COVER.
- MAKE SURE THAT YOU HAVE USED THE SAME SET OF AP NUMBER LABELS ON ALL AP EXAMS YOU HAVE TAKEN THIS YEAR.

Do not write beyond this border.

NO CALCULATOR ALLOWED

Work for problem 6(a)

No odd terms

$$P_{(x)} = 7 + (x-2) \overset{0}{\cancel{f'(2)}} + \frac{1}{9} \cdot (x-2)^2 \cdot \frac{1}{2!}$$

f(x)

$$\frac{6}{81}$$

$$P_8(x) = 7 + \frac{1!(x-2)^2}{9 \cdot 2!} + \frac{3!(x-2)^4}{3^4 \cdot 4!} + \frac{5!(x-2)^6}{3^6 \cdot 6!}$$

$$\frac{5!}{3^5} \quad P_8(x) = 7 + \frac{(x-2)^2}{9 \cdot 2} + \frac{(x-2)^4}{3^4 \cdot 4} + \frac{(x-2)^6}{3^6 \cdot 6}$$


Work for problem 6(b)

$$\frac{(x-2)^{2n}}{3^{2n} \cdot (2n)} \quad n \geq 1$$

Coefficient is  $\frac{1}{3^{2n} \cdot (2n)}$

Do not write beyond this border.

Continue problem 6 on page 15.



## NO CALCULATOR ALLOWED

Work for problem 6(c)

$$7 + \sum_{n=1}^{\infty} \frac{(x-2)^{2n}}{3^{2n} \cdot (2n)}$$

$$\lim_{n \rightarrow \infty} \left| \frac{(x-2)^{2n+2}}{3^{2n+2} \cdot (2n+2)} \cdot \frac{3^{2n} (2n)}{(x-2)^{2n}} \right| =$$

Left hand  $x=2$ 

$$\lim_{n \rightarrow \infty} = \left| \frac{(x-2)^2 (2n)}{9 (2n+2)} \right| < 1$$

$$\sum_{n=1}^{\infty} \frac{0}{3^{2n} \cdot (2n)} = 0$$

converges

Zinbolsive

Right hand  $x=5$ 

$$\sum_{n=1}^{\infty} \frac{(3)^{2n}}{3^{2n} \cdot (2n)} = \frac{1}{2n}$$

harmonic series diverges

$$-1 < \frac{(x-2)^2}{9} < 1$$

But same has to be 0

$$-9 < (x-2)^2 < 9$$

$$0 < x-2 < 3$$

$$2 < x < 5$$

Therefore  $\sum_{n=1}^{\infty} \frac{(x-2)^{2n}}{3^{2n} \cdot 2n}$  has interval of convergence of  $[2, 5)$

END OF EXAM

Do not write beyond this border.

THE FOLLOWING INSTRUCTIONS APPLY TO THE BACK COVER OF THIS SECTION II BOOKLET.

- MAKE SURE YOU HAVE COMPLETED THE IDENTIFICATION INFORMATION AS REQUESTED ON THE BACK OF THIS SECTION II BOOKLET.
- CHECK TO SEE THAT YOUR AP NUMBER APPEARS IN THE BOX(ES) ON THE BACK COVER.
- MAKE SURE THAT YOU HAVE USED THE SAME SET OF AP NUMBER LABELS ON ALL AP EXAMS YOU HAVE TAKEN THIS YEAR.

NO CALCULATOR ALLOWED

Work for problem 6(a)

$f(2) = 7.$

1st, 3rd, 5th derivatives = 0.

$f'(2) = 0$      $f''(2) = \frac{1}{9}$   
 $f'''(2) = 0$      $f^{(4)}(2) = \frac{3!}{3^4}$   
 $f^{(5)}(2) = 0$      $f^{(6)}(2) = \frac{5!}{3^6}$

Taylor Series

$$f(x) = f(x) + \frac{f'(x)(x-a)}{1!} + \frac{f''(x)(x-a)^2}{2!} + \frac{f'''(x)(x-a)^3}{3!} + \dots$$

$$7 + 0 + \frac{1}{9}(x-2)^2 + 0 + \frac{3!}{3^4}(x-2)^4 + 0 + \frac{5!}{3^6}(x-2)^6$$

$$= 7 + \frac{1}{18}(x-2)^2 + \frac{(x-2)^4}{3^4 \cdot 4} + \frac{(x-2)^6}{3^6 \cdot 6}$$

Work for problem 6(b)

general term for Taylor series about  $x = 2$

$$\frac{(x-2)^{2n}}{3^{2n} \cdot n}$$

The coefficient of  $(x-2)^{2n}$  for  $n \geq 1$  is 0 when  $n$  is odd and  $\frac{1}{3^{2n} \cdot n}$  when  $n$  is even and  $n \geq 2$ .

Do not write beyond this border.

## NO CALCULATOR ALLOWED

Work for problem 6(c)

Interval of convergence using Ratio test

$$\lim_{n \rightarrow \infty} \left| \frac{(x-2)^{(2n+2)} \cdot 3^{2n} \cdot n}{3^{2n+2} \cdot (n+1) \cdot (x-2)^{2n}} \right|$$

$$\lim_{n \rightarrow \infty} \left| \frac{(x-2)^2 n}{3^2 (n+1)} \right| = \lim_{n \rightarrow \infty} | (x-2)^2 |$$

\* as  $n \rightarrow \infty$ , the

END OF EXAM

THE FOLLOWING INSTRUCTIONS APPLY TO THE BACK COVER OF THIS SECTION II BOOKLET.

- MAKE SURE YOU HAVE COMPLETED THE IDENTIFICATION INFORMATION AS REQUESTED ON THE BACK OF THIS SECTION II BOOKLET.
- CHECK TO SEE THAT YOUR AP NUMBER APPEARS IN THE BOX(ES) ON THE BACK COVER.
- MAKE SURE THAT YOU HAVE USED THE SAME SET OF AP NUMBER LABELS ON ALL AP EXAMS YOU HAVE TAKEN THIS YEAR.

Do not write beyond this border.