



AP[®] Calculus BC 2003 Free-Response Questions Form B

The materials included in these files are intended for use by AP teachers for course and exam preparation; permission for any other use must be sought from the Advanced Placement Program[®]. Teachers may reproduce them, in whole or in part, in limited quantities for noncommercial, face-to-face teaching purposes. This permission does not apply to any third-party copyrights contained herein. This material may not be mass distributed, electronically or otherwise. These materials and any copies made of them may not be resold, and the copyright notices must be retained as they appear here.

These materials were produced by Educational Testing Service[®] (ETS[®]), which develops and administers the examinations of the Advanced Placement Program for the College Board. The College Board and Educational Testing Service (ETS) are dedicated to the principle of equal opportunity, and their programs, services, and employment policies are guided by that principle.

The College Board is a national nonprofit membership association whose mission is to prepare, inspire, and connect students to college and opportunity. Founded in 1900, the association is composed of more than 4,300 schools, colleges, universities, and other educational organizations. Each year, the College Board serves over three million students and their parents, 22,000 high schools, and 3,500 colleges through major programs and services in college admissions, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT[®], the PSAT/NMSQT[®], and the Advanced Placement Program[®] (AP[®]). The College Board is committed to the principles of equity and excellence, and that commitment is embodied in all of its programs, services, activities, and concerns.

For further information, visit www.collegeboard.com

Copyright © 2003 College Entrance Examination Board. All rights reserved. College Board, Advanced Placement Program, AP, AP Vertical Teams, APCD, Pacesetter, Pre-AP, SAT, Student Search Service, and the acorn logo are registered trademarks of the College Entrance Examination Board. AP Central is a trademark owned by the College Entrance Examination Board. PSAT/NMSQT is a registered trademark jointly owned by the College Entrance Examination Board and the National Merit Scholarship Corporation. Educational Testing Service and ETS are registered trademarks of Educational Testing Service. Other products and services may be trademarks of their respective owners.

For the College Board's online home for AP professionals, visit AP Central at apcentral.collegeboard.com.


2003 AP[®] CALCULUS BC FREE-RESPONSE QUESTIONS (Form B)

CALCULUS BC
SECTION II, Part A

Time—45 minutes


Number of problems—3

A graphing calculator is required for some problems or parts of problems.



1. Let f be the function given by $f(x) = 4x^2 - x^3$, and let l be the line $y = 18 - 3x$, where l is tangent to the graph of f . Let R be the region bounded by the graph of f and the x -axis, and let S be the region bounded by the graph of f , the line l , and the x -axis, as shown above.
- (a) Show that l is tangent to the graph of $y = f(x)$ at the point $x = 3$.
- (b) Find the area of S .
- (c) Find the volume of the solid generated when R is revolved about the x -axis.
-

2003 AP[®] CALCULUS BC FREE-RESPONSE QUESTIONS (Form B)



2. The figure above shows the graphs of the circles $x^2 + y^2 = 2$ and $(x - 1)^2 + y^2 = 1$. The graphs intersect at the points $(1, 1)$ and $(1, -1)$. Let R be the shaded region in the first quadrant bounded by the two circles and the x -axis.
- Set up an expression involving one or more integrals with respect to x that represents the area of R .
 - Set up an expression involving one or more integrals with respect to y that represents the area of R .
 - The polar equations of the circles are $r = \sqrt{2}$ and $r = 2 \cos \theta$, respectively. Set up an expression involving one or more integrals with respect to the polar angle θ that represents the area of R .

Distance x (mm)	0	60	120	180	240	300	360
Diameter $B(x)$ (mm)	24	30	28	30	26	24	26

3. A blood vessel is 360 millimeters (mm) long with circular cross sections of varying diameter. The table above gives the measurements of the diameter of the blood vessel at selected points along the length of the blood vessel, where x represents the distance from one end of the blood vessel and $B(x)$ is a twice-differentiable function that represents the diameter at that point.
- Write an integral expression in terms of $B(x)$ that represents the average radius, in mm, of the blood vessel between $x = 0$ and $x = 360$.
 - Approximate the value of your answer from part (a) using the data from the table and a midpoint Riemann sum with three subintervals of equal length. Show the computations that lead to your answer.
 - Using correct units, explain the meaning of $\pi \int_{125}^{275} \left(\frac{B(x)}{2} \right)^2 dx$ in terms of the blood vessel.
 - Explain why there must be at least one value x , for $0 < x < 360$, such that $B''(x) = 0$.

END OF PART A OF SECTION II

Copyright © 2003 by College Entrance Examination Board. All rights reserved.
 Available to AP professionals at apcentral.collegeboard.com and to
 students and parents at www.collegeboard.com/apstudents.

2003 AP[®] CALCULUS BC FREE-RESPONSE QUESTIONS (Form B)

CALCULUS BC
SECTION II, Part B

Time—45 minutes


Number of problems—3

No calculator is allowed for these problems.

4. A particle moves in the xy -plane so that the position of the particle at any time t is given by

$$x(t) = 2e^{3t} + e^{-7t} \text{ and } y(t) = 3e^{3t} - e^{-2t}.$$

- (a) Find the velocity vector for the particle in terms of t , and find the speed of the particle at time $t = 0$.
- (b) Find $\frac{dy}{dx}$ in terms of t , and find $\lim_{t \rightarrow \infty} \frac{dy}{dx}$.
- (c) Find each value t at which the line tangent to the path of the particle is horizontal, or explain why none exists.
- (d) Find each value t at which the line tangent to the path of the particle is vertical, or explain why none exists.



Graph of f

5. Let f be a function defined on the closed interval $[0, 7]$. The graph of f , consisting of four line segments, is shown above. Let g be the function given by $g(x) = \int_2^x f(t) dt$.
- (a) Find $g(3)$, $g'(3)$, and $g''(3)$.
- (b) Find the average rate of change of g on the interval $0 \leq x \leq 3$.
- (c) For how many values c , where $0 < c < 3$, is $g'(c)$ equal to the average rate found in part (b)? Explain your reasoning.
- (d) Find the x -coordinate of each point of inflection of the graph of g on the interval $0 < x < 7$. Justify your answer.

2003 AP[®] CALCULUS BC FREE-RESPONSE QUESTIONS (Form B)

6. The function f has a Taylor series about $x = 2$ that converges to $f(x)$ for all x in the interval of convergence. The n th derivative of f at $x = 2$ is given by $f^{(n)}(2) = \frac{(n+1)!}{3^n}$ for $n \geq 1$, and $f(2) = 1$.
- (a) Write the first four terms and the general term of the Taylor series for f about $x = 2$.
 - (b) Find the radius of convergence for the Taylor series for f about $x = 2$. Show the work that leads to your answer.
 - (c) Let g be a function satisfying $g(2) = 3$ and $g'(x) = f(x)$ for all x . Write the first four terms and the general term of the Taylor series for g about $x = 2$.
 - (d) Does the Taylor series for g as defined in part (c) converge at $x = -2$? Give a reason for your answer.
-

END OF EXAMINATION