



**AP<sup>®</sup>**  
**Calculus BC**  
1998 Scoring  
Guidelines

connect to college success™  
[www.collegeboard.com](http://www.collegeboard.com)

## The College Board: Connecting Students to College Success

The College Board is a not-for-profit membership association whose mission is to connect students to college success and opportunity. Founded in 1900, the association is composed of more than 4,700 schools, colleges, universities, and other educational organizations. Each year, the College Board serves over three and a half million students and their parents, 23,000 high schools, and 3,500 colleges through major programs and services in college admissions, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT<sup>®</sup>, the PSAT/NMSQT<sup>®</sup>, and the Advanced Placement Program<sup>®</sup> (AP<sup>®</sup>). The College Board is committed to the principles of excellence and equity, and that commitment is embodied in all of its programs, services, activities, and concerns.

### Permission to Reprint Statement

The College Board intends this publication for noncommercial use by teachers for course and exam preparation; permission for any other use must be sought from the College Board. Teachers may reproduce this publication, in whole or in part, **in limited print quantities for noncommercial, face-to-face teaching purposes** and distribute up to 50 print copies from a teacher to a class of middle or high school students, with each student receiving no more than one copy.

This permission does not apply to any third-party copyrights contained within this publication.

When educators reproduce this publication for noncommercial, face-to-face teaching purposes, the following source line must be included:

**“AP Calculus BC, 1998 Scoring Guidelines.” Copyright © 1998 by the College Board. Reprinted with permission. All rights reserved. [www.collegeboard.com](http://www.collegeboard.com). This material may not be mass distributed, electronically or otherwise. This publication and any copies made from it may not be resold.**

No party may share this copyrighted material electronically—by fax, Web site, CD-ROM, disk, e-mail, electronic discussion group, or any other electronic means not stated here. In some cases—such as online courses or online workshops—the College Board *may* grant permission for electronic dissemination of its copyrighted materials. All intended uses not defined within ***noncommercial, face-to-face teaching purposes*** (including distribution exceeding 50 copies) must be reviewed and approved; in these cases, a license agreement must be received and signed by the requestor and copyright owners prior to the use of copyrighted material. Depending on the nature of the request, a licensing fee may be applied. Please use the required form accessible online. The form may be found at:  
[www.collegeboard.com/inquiry/cbpermit.html](http://www.collegeboard.com/inquiry/cbpermit.html).

## Equity Policy Statement

The College Board and the Advanced Placement Program encourage teachers, AP Coordinators, and school administrators to make equitable access a guiding principle for their AP programs. The College Board is committed to the principle that all students deserve an opportunity to participate in rigorous and academically challenging courses and programs. All students who are willing to accept the challenge of a rigorous academic curriculum should be considered for admission to AP courses. The Board encourages the elimination of barriers that restrict access to AP courses for students from ethnic, racial, and socioeconomic groups that have been traditionally underrepresented in the AP Program. Schools should make every effort to ensure that their AP classes reflect the diversity of their student population.

For more information about equity and access in principle and practice, please send an email to [apecquity@collegeboard.org](mailto:apecquity@collegeboard.org).


Copyright © 1998 by College Board. All rights reserved. College Board, AP Central, APCD, Advanced Placement Program, AP, AP Vertical Teams, Pre-AP, SAT, and the acorn logo are registered trademarks of the College Entrance Examination Board. Admitted Class Evaluation Service, CollegeEd, Connect to college success, MyRoad, SAT Professional Development, SAT Readiness Program, and Setting the Cornerstones are trademarks owned by the College Entrance Examination Board. PSAT/NMSQT is a trademark of the College Entrance Examination Board and National Merit Scholarship Corporation. Other products and services may be trademarks of their respective owners. Visit College Board on the Web: [www.collegeboard.com](http://www.collegeboard.com).

**For further information, visit [apcentral.collegeboard.com](http://apcentral.collegeboard.com).**

## 1998 Calculus BC Scoring Guidelines

1. Let  $R$  be the region in the first quadrant bounded by the graph of  $y = 8 - x^{3/2}$ , the  $x$ -axis, and the  $y$ -axis.
- (a) Find the area of the region  $R$ .
- (b) Find the volume of the solid generated when  $R$  is revolved about the  $x$ -axis.
- (c) The vertical line  $x = k$  divides the region  $R$  into two regions such that when these two regions are revolved about the  $x$ -axis, they generate solids with equal volumes. Find the value of  $k$ .

(a)



$$A = \int_0^4 (8 - x^{3/2}) dx$$

$$= 8x - \frac{2}{5}x^{5/2} \Big|_0^4 = 32 - \frac{64}{5} = \frac{96}{5} = 19.2$$

(b)  $V = \pi \int_0^4 (8 - x^{3/2})^2 dx$

$$= \frac{576\pi}{5} = 115.2\pi \approx 361.911$$

(c)  $\pi \int_0^k (8 - x^{3/2})^2 dx = \frac{115.2\pi}{2}$

[or

$$\left[ \pi \int_0^k (8 - x^{3/2})^2 dx = \pi \int_k^4 (8 - x^{3/2})^2 dx \right]$$

$$\int_0^k (8 - x^{3/2})^2 dx = 57.6$$

$$\int_0^k (64 - 16x^{3/2} + x^3) dx = 57.6$$

$$64k - \frac{32}{5}k^{5/2} + \frac{k^4}{4} = 57.6$$

$$k \approx 0.995 \text{ or } 0.994$$

3 {

- 2: integral
- 1: integrand
- 1: limits
- 1: answer

3 {

- 2: integral
- 1: integrand
- 1: limits and constant
- 1: answer

3 {

- 1: integral with  $k$  in limits
- 1: equates volumes
- 1: answer

Note: 0/1 for answer in each part if no setup points earned

## 1998 Calculus BC Scoring Guidelines

2. Let  $f$  be the function given by  $f(x) = 2xe^{2x}$ .
- (a) Find  $\lim_{x \rightarrow -\infty} f(x)$  and  $\lim_{x \rightarrow \infty} f(x)$ .
- (b) Find the absolute minimum value of  $f$ . Justify that your answer is an absolute minimum.
- (c) What is the range of  $f$ ?
- (d) Consider the family of functions defined by  $y = bxe^{bx}$ , where  $b$  is a nonzero constant. Show that the absolute minimum value of  $bxe^{bx}$  is the same for all nonzero values of  $b$ .

(a)  $\lim_{x \rightarrow -\infty} 2xe^{2x} = 0$

$\lim_{x \rightarrow \infty} 2xe^{2x} = \infty$  or DNE

(b)  $f'(x) = 2e^{2x} + 2x \cdot 2 \cdot e^{2x} = 2e^{2x}(1 + 2x) = 0$

if  $x = -1/2$

$f(-1/2) = -1/e$  or  $-0.368$  or  $-0.367$

$-1/e$  is an absolute minimum value because:

(i)  $f'(x) < 0$  for all  $x < -1/2$  and

$f'(x) > 0$  for all  $x > -1/2$

-or-

(ii)  $f'(x) \begin{array}{c} - \qquad \qquad + \\ \hline \qquad \qquad -1/2 \end{array}$

and  $x = -1/2$  is the only critical number

(c) Range of  $f = [-1/e, \infty)$

or  $[-0.367, \infty)$

or  $[-0.368, \infty)$

(d)  $y' = be^{bx} + b^2xe^{bx} = be^{bx}(1 + bx) = 0$

if  $x = -1/b$

At  $x = -1/b$ ,  $y = -1/e$

$y$  has an absolute minimum value of  $-1/e$  for all nonzero  $b$

2  $\left\{ \begin{array}{l} 1: 0 \text{ as } x \rightarrow -\infty \\ 1: \infty \text{ or DNE as } x \rightarrow \infty \end{array} \right.$

3  $\left\{ \begin{array}{l} 1: \text{solves } f'(x) = 0 \\ 1: \text{evaluates } f \text{ at student's critical point} \\ \quad 0/1 \text{ if not local minimum from} \\ \quad \text{student's derivative} \\ 1: \text{justifies absolute minimum value} \\ \quad 0/1 \text{ for a local argument} \\ \quad 0/1 \text{ without explicit symbolic} \\ \quad \text{derivative} \end{array} \right.$

Note: 0/3 if no absolute minimum based on student's derivative

1: answer

Note: must include the left-hand endpoint; exclude the right-hand "endpoint"

3  $\left\{ \begin{array}{l} 1: \text{sets } y' = be^{bx}(1 + bx) = 0 \\ 1: \text{solves student's } y' = 0 \\ 1: \text{evaluates } y \text{ at a critical number} \\ \quad \text{and gets a value independent of } b \end{array} \right.$

Note: 0/3 if only considering specific values of  $b$

## 1998 Calculus BC Scoring Guidelines

3. Let  $f$  be a function that has derivatives of all orders for all real numbers. Assume  $f(0) = 5$ ,  $f'(0) = -3$ ,  $f''(0) = 1$ , and  $f'''(0) = 4$ .

- (a) Write the third-degree Taylor polynomial for  $f$  about  $x = 0$  and use it to approximate  $f(0.2)$ .
- (b) Write the fourth-degree Taylor polynomial for  $g$ , where  $g(x) = f(x^2)$ , about  $x = 0$ .
- (c) Write the third-degree Taylor polynomial for  $h$ , where  $h(x) = \int_0^x f(t) dt$ , about  $x = 0$ .
- (d) Let  $h$  be defined as in part (c). Given that  $f(1) = 3$ , either find the exact value of  $h(1)$  or explain why it cannot be determined.

(a) 
$$P_3(f)(x) = 5 - 3x + \frac{1}{2}x^2 + \frac{2}{3}x^3$$

$$f(0.2) \approx P_3(f)(0.2) =$$

$$5 - 3(0.2) + \frac{0.04}{2} + \frac{2(0.008)}{3} =$$

$$4.425$$

$$3 \left\{ \begin{array}{l} 2: 5 - 3x + \frac{1}{2}x^2 + \frac{2}{3}x^3 \\ <-1> \text{ each incorrect term,} \\ & \text{extra term, or } + \dots \\ 1: \text{ approximates } f(0.2) \end{array} \right.$$

<-1> for incorrect use of =

(b) 
$$P_4(g)(x) = P_2(f)(x^2) = 5 - 3x^2 + \frac{1}{2}x^4$$

2:  $P_2(f)(x^2)$   
<-1> each incorrect or extra term

(c) 
$$P_3(h)(x) = \int_0^x \left( 5 - 3t + \frac{1}{2}t^2 \right) dt$$

$$= \left[ 5t - \frac{3}{2}t^2 + \frac{1}{6}t^3 \right]_0^x$$

$$= 5x - \frac{3}{2}x^2 + \frac{1}{6}x^3$$

$$2 \left\{ \begin{array}{l} 1: P_3(h)(x) = \int_0^x P_2(f)(t) dt \\ 1: \text{ answer} \\ 0/1 \text{ if any incorrect or extra terms} \end{array} \right.$$

(d) 
$$h(1) = \int_0^1 f(t) dt$$


cannot be determined because  $f(t)$  is known only for  $t = 0$  and  $t = 1$

$$2 \left\{ \begin{array}{l} 1: h(1) \text{ cannot be determined} \\ 1: \text{ reason} \end{array} \right.$$

## 1998 Calculus BC Scoring Guidelines

4. Consider the differential equation given by  $\frac{dy}{dx} = \frac{xy}{2}$ .
- (a) On the axes provided below, sketch a slope field for the given differential equation at the nine points indicated.
- (b) Let  $y = f(x)$  be the particular solution to the given differential equation with the initial condition  $f(0) = 3$ . Use Euler's method starting at  $x = 0$ , with a step size of 0.1, to approximate  $f(0.2)$ . Show the work that leads to your answer.
- (c) Find the particular solution  $y = f(x)$  to the given differential equation with the initial condition  $f(0) = 3$ . Use your solution to find  $f(0.2)$ .

(a)



(b)  $f(0.1) \approx f(0) + f'(0)(0.1)$   
 $= 3 + \frac{1}{2}(0)(3)(0.1) = 3$   
 $f(0.2) \approx f(0.1) + f'(0.1)(0.1)$   
 $= 3 + \frac{1}{2}(0.1)(3)(0.1)$   
 $= 3 + \frac{.03}{2} = 3.015$

(c)  $\frac{dy}{dx} = \frac{xy}{2}$   
 $\int \frac{dy}{y} = \int \frac{x}{2} dx$   
 $\ln |y| = \frac{1}{4}x^2 + C_1$   
 $y = Ce^{x^2/4}$   
 $3 = Ce^0 \implies C = 3$   
 $y = 3e^{x^2/4}$   
 $f(0.2) = 3e^{.04/4} = 3e^{.01} = 3.030$

1: line segments at nine points with negative - zero - positive slope left to right and increasing steepness bottom to top at  $x = 1$  and  $x = -1$

2 { 1: Euler's Method equations or equivalent table  
 1: answer (not eligible without first point)

Special Case: 1/2 for first iteration 3.015 and second iteration 3.045

6 { 1: separates variables  
 1: antiderivative of  $dy$  term  
 1: antiderivative of  $dx$  term  
 1: solves for  $y$   
 1: solves for constant of integration  
 1: evaluates  $f(0.2)$

Note: max 4/6 [1-1-1-0-0-1] if no constant of integration


## 1998 Calculus BC Scoring Guidelines

5. The temperature outside a house during a 24-hour period is given by

$$F(t) = 80 - 10 \cos\left(\frac{\pi t}{12}\right), \quad 0 \leq t \leq 24,$$

where  $F(t)$  is measured in degrees Fahrenheit and  $t$  is measured in hours.

- Sketch the graph of  $F$  on the grid below.
- Find the average temperature, to the nearest degree Fahrenheit, between  $t = 6$  and  $t = 14$ .
- An air conditioner cooled the house whenever the outside temperature was at or above 78 degrees Fahrenheit. For what values of  $t$  was the air conditioner cooling the house?
- The cost of cooling the house accumulates at the rate of \$0.05 per hour for each degree the outside temperature exceeds 78 degrees Fahrenheit. What was the total cost, to the nearest cent, to cool the house for this 24-hour period?



(b)

$$\begin{aligned} \text{Avg.} &= \frac{1}{14-6} \int_6^{14} \left[ 80 - 10 \cos\left(\frac{\pi t}{12}\right) \right] dt \\ &= \frac{1}{8} (697.2957795) \\ &= 87.162 \text{ or } 87.161 \\ &\approx 87^\circ \text{ F} \end{aligned}$$

(c)

$$\begin{aligned} \left[ 80 - 10 \cos\left(\frac{\pi t}{12}\right) \right] - 78 &\geq 0 \\ 2 - 10 \cos\left(\frac{\pi t}{12}\right) &\geq 0 \\ \left. \begin{array}{l} 5.230 \\ \text{or} \\ 5.231 \end{array} \right\} &\leq t \leq \left\{ \begin{array}{l} 18.769 \\ \text{or} \\ 18.770 \end{array} \right. \end{aligned}$$

(d)

$$\begin{aligned} C &= 0.05 \int_{\substack{5.231 \\ \text{or} \\ 5.230}}^{\substack{18.769 \\ \text{or} \\ 18.770}} \left( \left[ 80 - 10 \cos\left(\frac{\pi t}{12}\right) \right] - 78 \right) dt \\ &= 0.05(101.92741) = 5.096 \approx \$5.10 \end{aligned}$$

1: bell-shaped graph  
 minimum 70 at  $t = 0, t = 24$  only  
 maximum 90 at  $t = 12$  only

2: integral  
 1: limits and  $1/(14-6)$   
 1: integrand  
 3 { 1: answer  
 0/1 if integral not of the form  
 $\frac{1}{b-a} \int_a^b F(t) dt$

2 { 1: inequality or equation  
 1: solutions with interval

2: integral  
 1: limits and 0.05  
 1: integrand  
 3 { 1: answer  
 0/1 if integral not of the form  
 $k \int_a^b (F(t) - 78) dt$

## 1998 Calculus BC Scoring Guidelines

6. A particle moves along the curve defined by the equation  $y = x^3 - 3x$ . The  $x$ -coordinate of the particle,  $x(t)$ , satisfies the equation  $\frac{dx}{dt} = \frac{1}{\sqrt{2t+1}}$ , for  $t \geq 0$  with initial condition  $x(0) = -4$ .
- (a) Find  $x(t)$  in terms of  $t$ .
- (b) Find  $\frac{dy}{dt}$  in terms of  $t$ .
- (c) Find the location and speed of the particle at time  $t = 4$ .

(a)  $x(t) = \int \frac{1}{\sqrt{2t+1}} dt$   
 $x(t) = \sqrt{2t+1} + C$   
 $x(0) = -4 = 1 + C \implies C = -5$   
 $x(t) = \sqrt{2t+1} - 5$

(b)  $y = x^3 - 3x$   
 $\frac{dy}{dt} = 3x^2 \frac{dx}{dt} - 3 \frac{dx}{dt}$   
 $= (3x^2 - 3) \frac{dx}{dt}$   
 $= \left[ 3(\sqrt{2t+1} - 5)^2 - 3 \right] \left[ \frac{1}{\sqrt{2t+1}} \right]$

(c)  $x(4) = \sqrt{9} - 5 = -2$   
 $y(4) = (-2)^3 - 3(-2) = -2$   
 Location at  $t = 4$  is  $(-2, -2)$   
 $\left. \frac{dx}{dt} \right|_{t=4} = \frac{1}{3}$

$$\left. \frac{dy}{dt} \right|_{t=4} = \frac{3(3-5)^2 - 3}{3} = 3$$

$$\text{Speed} = \sqrt{\left(\frac{1}{3}\right)^2 + 3^2} = \sqrt{\frac{82}{9}} = 3.018$$

3  $\left\{ \begin{array}{l} 1: x(t) = \int \frac{dt}{\sqrt{2t+1}} \\ 1: x(t) = \sqrt{2t+1} + C \\ 1: \text{evaluates } C \end{array} \right.$

2: answer

<-1> each error

Note: failure to express  $\frac{dy}{dt}$  solely in terms of  $t$  is a single error

4  $\left\{ \begin{array}{l} 1: \text{position} \\ 1: \text{evaluates } \frac{dx}{dt} \text{ and } \frac{dy}{dt} \text{ at } t = 4 \\ 1: \text{uses speed formula} \\ 1: \text{answer} \end{array} \right.$