




## AP<sup>®</sup> Physics C: Mechanics 2001 Sample Student Responses

**The materials included in these files are intended for non-commercial use by AP teachers for course and exam preparation; permission for any other use must be sought from the Advanced Placement Program. Teachers may reproduce them, in whole or in part, in limited quantities, for face-to-face teaching purposes but may not mass distribute the materials, electronically or otherwise. These materials and any copies made of them may not be resold, and the copyright notices must be retained as they appear here. This permission does not apply to any third-party copyrights contained herein.**

These materials were produced by Educational Testing Service (ETS), which develops and administers the examinations of the Advanced Placement Program for the College Board. The College Board and Educational Testing Service (ETS) are dedicated to the principle of equal opportunity, and their programs, services, and employment policies are guided by that principle.

The College Board is a national nonprofit membership association dedicated to preparing, inspiring, and connecting students to college and opportunity. Founded in 1900, the association is composed of more than 3,900 schools, colleges, universities, and other educational organizations. Each year, the College Board serves over three million students and their parents, 22,000 high schools, and 3,500 colleges, through major programs and services in college admission, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT<sup>®</sup>, the PSAT/NMSQT<sup>™</sup>, the Advanced Placement Program<sup>®</sup> (AP<sup>®</sup>), and Pacesetter<sup>®</sup>. The College Board is committed to the principles of equity and excellence, and that commitment is embodied in all of its programs, services, activities, and concerns.

Copyright © 2001 by College Entrance Examination Board. All rights reserved. College Board, Advanced Placement Program, AP, and the acorn logo are registered trademarks of the College Entrance Examination Board.



Experiment A

Mech 3.

A light string that is attached to a large block of mass  $4m$  passes over a pulley with negligible rotational inertia and is wrapped around a vertical pole of radius  $r$ , as shown in Experiment A above. The system is released from rest, and as the block descends the string unwinds and the vertical pole with its attached apparatus rotates. The apparatus consists of a horizontal rod of length  $2L$ , with a small block of mass  $m$  attached at each end. The rotational inertia of the pole and the rod are negligible.


(a) Determine the rotational inertia of the rod-and-block apparatus attached to the top of the pole.

$$I = \sum mr^2$$

$$I = mL^2 + mL^2$$

$$I = 2mL^2$$

(b) Determine the downward acceleration of the large block.



$$\Sigma F_x = \vec{w} + \vec{T} = 4ma$$

$$4mg - T = 4ma$$

$$\Sigma \tau = \tau_T = I\alpha$$

$$\vec{r} \times \vec{T} = 2mL^2 \frac{a}{r}$$

$$rT = 2mL^2 \frac{a}{r}$$

$$T = \frac{2mL^2 a}{r^2}$$

$$4mg - \frac{2mL^2 a}{r^2} = 4ma$$

$$4mg = a \left( 4m + \frac{2mL^2}{r^2} \right)$$

$$2g = a \left( 2 + \frac{L^2}{r^2} \right)$$

$$\frac{2g}{2 + \frac{L^2}{r^2}} = a$$


GO ON TO THE NEXT PAGE.

(c) When the large block has descended a distance  $D$ , how does the instantaneous total kinetic energy of the three blocks compare with the value  $4mgD$ ? Check the appropriate space below.

\_\_\_ Greater than  $4mgD$     X Equal to  $4mgD$     \_\_\_ Less than  $4mgD$

Justify your answer.

Mechanical Energy is conserved, so  
 $\Delta K = -\Delta U$   
 $K_{tot} = 4mgD$



Experiment B

The system is now reset. The string is rewound around the pole to bring the large block back to its original location. The small blocks are detached from the rod and then suspended from each end of the rod, using strings of length  $l$ . The system is again released from rest so that as the large block descends and the apparatus rotates, the small blocks swing outward, as shown in Experiment B above. This time the downward acceleration of the block decreases with time after the system is released.


(d) When the large block has descended a distance  $D$ , how does the instantaneous total kinetic energy of the three blocks compare to that in part (c)? Check the appropriate space below.

\_\_\_ Greater    \_\_\_ Equal    X Less

Justify your answer.


Mechanical energy is still conserved, but some of that energy has gone into lifting the blocks on the rod (as potential energy),

like this:



so the total kinetic energy is less than before.

GO ON TO THE NEXT PAGE.



Experiment A

Mech 3.

A light string that is attached to a large block of mass  $4m$  passes over a pulley with negligible rotational inertia and is wrapped around a vertical pole of radius  $r$ , as shown in Experiment A above. The system is released from rest, and as the block descends the string unwinds and the vertical pole with its attached apparatus rotates. The apparatus consists of a horizontal rod of length  $2L$ , with a small block of mass  $m$  attached at each end. The rotational inertia of the pole and the rod are negligible.

(a) Determine the rotational inertia of the rod-and-block apparatus attached to the top of the pole.

$I = M L^2 + M L^2$  . *rod rotational inertia is negligible*

$I = 2ML^2$

(b) Determine the downward acceleration of the large block.

$\Sigma F = 4ma$

$4ma = 4mg - T$

$Ia = rT$

$I \frac{a}{r} = rT$

$2ML^2 \frac{a}{r} = rT$

$4Ma = 4Mg - \frac{2ML^2 a}{r^2}$

$4Ma + \frac{2ML^2 a}{r^2} = 4Mg$

$a \left( 4M + \frac{2ML^2}{r^2} \right) = 4Mg$

$a = \frac{4g}{4 + \frac{2L^2}{r^2}}$

$a = \frac{g}{1 + \frac{L^2}{2r^2}}$

$\Sigma \tau = I \alpha$

$\frac{2ML^2 a}{r^2} = +$

GO ON TO THE NEXT PAGE.


(c) When the large block has descended a distance  $D$ , how does the instantaneous total kinetic energy of the three blocks compare with the value  $4mgD$ ? Check the appropriate space below.

\_\_\_ Greater than  $4mgD$     X Equal to  $4mgD$     \_\_\_ Less than  $4mgD$

Justify your answer.

No  
initial  
 $K_i$

$K_1 + U_{g1} = K_2 + U_{g2}$  ← No initial potential energy in Block  
 $U_{g1} = K_2$   
 $4mgd = \frac{1}{2} I \omega^2 + \frac{1}{2} m \omega^2$     SO.  $4mgd = \text{total } K_e$



Experiment B

The system is now reset. The string is rewound around the pole to bring the large block back to its original location. The small blocks are detached from the rod and then suspended from each end of the rod, using strings of length  $l$ . The system is again released from rest so that as the large block descends and the apparatus rotates, the small blocks swing outward, as shown in Experiment B above. This time the downward acceleration of the block decreases with time after the system is released.

(d) When the large block has descended a distance  $D$ , how does the instantaneous total kinetic energy of the three blocks compare to that in part (c)? Check the appropriate space below.

\_\_\_ Greater    X Equal    \_\_\_ Less

Justify your answer.

No energy is being lost. As the blocks swing out they simply increase the rotational inertia of the rod and blocks, this accounts for the decreasing acceleration. But no energy is lost to friction or other forces.

$K_1 + U_{g1} = K_2 + U_{g2}$   
 Still  $U_{g1} = K_2$  = no energy lost    GO ON TO THE NEXT PAGE.  
 -9- total  $K_e$  is same.