



## AP Physics C: Electricity & Magnetism 1999 Scoring Guidelines

**The materials included in these files are intended for non-commercial use by AP teachers for course and exam preparation; permission for any other use must be sought from the Advanced Placement Program. Teachers may reproduce them, in whole or in part, in limited quantities, for face-to-face teaching purposes but may not mass distribute the materials, electronically or otherwise. These materials and any copies made of them may not be resold, and the copyright notices must be retained as they appear here. This permission does not apply to any third-party copyrights contained herein.**

These materials were produced by Educational Testing Service (ETS), which develops and administers the examinations of the Advanced Placement Program for the College Board. The College Board and Educational Testing Service (ETS) are dedicated to the principle of equal opportunity, and their programs, services, and employment policies are guided by that principle.

The College Board is a national nonprofit membership association dedicated to preparing, inspiring, and connecting students to college and opportunity. Founded in 1900, the association is composed of more than 3,900 schools, colleges, universities, and other educational organizations. Each year, the College Board serves over three million students and their parents, 22,000 high schools, and 3,500 colleges, through major programs and services in college admission, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT<sup>®</sup>, the PSAT/NMSQT<sup>™</sup>, the Advanced Placement Program<sup>®</sup> (AP<sup>®</sup>), and Pacesetter<sup>®</sup>. The College Board is committed to the principles of equity and excellence, and that commitment is embodied in all of its programs, services, activities, and concerns.

Copyright © 2001 by College Entrance Examination Board. All rights reserved. College Board, Advanced Placement Program, AP, and the acorn logo are registered trademarks of the College Entrance Examination Board.

E &amp; M 1 (15 points)

(a) 4 points

For using the relationship between potential and charge

1 point

$$V = \frac{1}{4\pi\epsilon_0} \frac{Q}{r}$$

Solving for  $Q$ :

$$Q = 4\pi\epsilon_0 Vr$$

For correct substitutions for the potential and radius

1 point

$$Q_0 = 4\pi\epsilon_0 (-2000 \text{ V})(0.20 \text{ m}) \quad \text{or} \quad (-2000 \text{ V})(0.20 \text{ m}) / (9 \times 10^9 \text{ N}\cdot\text{m}^2/\text{C}^2)$$

$$Q_0 = -1600\pi\epsilon_0 \text{ C} \quad \text{or} \quad -4.4 \times 10^{-4} \text{ C}$$

For the correct magnitude of  $Q_0$ 

1 point

For the negative sign

1 point

(b) 5 points

i. For indicating that the electric field is zero

1 point

ii. The charge on the sphere can be treated as a point charge at its center

$$E = \frac{1}{4\pi\epsilon_0} \frac{Q_0}{r^2}$$

$$E = (9 \times 10^9 \text{ N}\cdot\text{m}^2/\text{C}^2) \left( \frac{4.4 \times 10^{-6} \text{ C}}{r^2} \right)$$

$$E = \frac{396 \text{ N}}{r^2} \quad \text{or} \quad \frac{400 \text{ N}}{r^2} \quad \text{where } r \text{ is in meters}$$

For any of the above expressions for  $E$ 

1 point

iii. For indicating that the electric field is zero

1 point

iv. For indicating that the electric field is zero

1 point

For having all four answers correct OR for some mention of using the enclosed charge OR for some mention of Gauss' law

1 point

E &amp; M 1 (continued)

(c) 3 points

$$\Delta V = V_b - V_a = - \int_a^b E \, dr$$

For recognition of the need to take the difference of the potentials at radii  $a$  and  $b$ ,  
or for writing the definite integral (with limits)

1 point

$$\begin{aligned} |\Delta V| &= \frac{Q_0}{4\pi\epsilon_0} \int_a^b \frac{dr}{r^2} \\ &= \frac{Q_0}{4\pi\epsilon_0} \left( \frac{1}{r} \right) \Big|_a^b \end{aligned}$$

$$|\Delta V| = \frac{Q_0}{4\pi\epsilon_0} \left( \frac{1}{b} - \frac{1}{a} \right)$$

For correct substitution of variables or numerical values for  $Q_0$ ,  $a$ , and  $b$

1 point

For the correct answer

1 point

$$|\Delta V| = \frac{5Q_0}{8\pi\epsilon_0} \quad \text{or} \quad 1000 \text{ V}$$

*(Alternate solution)**(Alternate points)*

For recognition of the need to take the difference of the potentials at radii  $a$  and  $b$

1 point

$$\Delta V = V_b - V_a$$

$$\Delta V = \frac{Q_0}{4\pi\epsilon_0} \left( \frac{1}{r_b} \right) - \frac{Q_0}{4\pi\epsilon_0} \left( \frac{1}{r_a} \right)$$

For correct substitution of  $Q_0$ ,  $a$ , and  $b$

1 point

$$\Delta V = \frac{Q_0}{4\pi\epsilon_0} \left( \frac{1}{b} - \frac{1}{a} \right)$$

For the correct answer

1 point

$$|\Delta V| = \frac{5Q_0}{8\pi\epsilon_0} \quad \text{or} \quad 1000 \text{ V}$$

*(Alternate solution)**(Alternate points)*

$$V = \frac{Q}{C}$$

For using the above relationship

1 point

For substituting  $Q_0$  from part (a) and  $C$  from part (d) alternate solution

1 point

For the correct answer

1 point

$$|\Delta V| = \frac{5Q_0}{8\pi\epsilon_0} \quad \text{or} \quad 1000 \text{ V}$$

E &amp; M 1 (continued)

(d) 2 points

$$C = \frac{Q_0}{V}$$

For using the above relationship

1 point

For substituting  $Q_0$  from part (a) and  $\Delta V$  from part (c)

1 point

$$C = \frac{4.4 \times 10^{-8} \text{ C}}{1000 \text{ V}}$$

$$C = 4.4 \times 10^{-11} \text{ F}$$

*(Alternate solution)**(Alternate points)*

For writing the equation for the capacitance of the spherical capacitor

1 point

$$C = \frac{4\pi\epsilon_0 ab}{b-a}$$

$$C = \frac{(0.02 \text{ m})(0.04 \text{ m})}{(9 \times 10^9 \text{ N}\cdot\text{m}^2/\text{C}^2)(0.04 \text{ m} - 0.02 \text{ m})}$$

For the correct answer

1 point

$$C = 4.4 \times 10^{-11} \text{ F}$$

For correct units on two answers and no incorrect units

1 point

## E &amp; M 2 (15 points)

## (a) 5 points

For using Faraday's law for a loop

1 point

$$\mathcal{E} = -\frac{d\phi}{dt} \quad \text{or} \quad \mathcal{E} = -\frac{\Delta\phi}{\Delta t}$$

For relating magnetic flux to magnetic field and area

1 point

$$\frac{d\phi}{dt} = A \frac{dB}{dt} \quad \text{or} \quad \frac{\Delta\phi}{\Delta t} = A \frac{\Delta B}{\Delta t}$$

For using the proper expression for the area of a loop

1 point

$$A = \pi r^2$$

$$\mathcal{E} = \pi r^2 \frac{dB}{dt} \quad \text{or} \quad \mathcal{E} = \pi r^2 \frac{\Delta B}{\Delta t}$$

For using the correct radius, i.e. the radius of the field

1 point

$$\mathcal{E} = \pi (0.6 \text{ m})^2 (0.40 \text{ T/s})$$

For computing the correct answer

1 point

$$\mathcal{E} = 0.45 \text{ V}$$

## (b) 3 points

For any statement of Ohm's law

1 point

$$V = IR$$

Solving for the current:

$$I = V/R = \mathcal{E}/R$$

$$I = (0.45 \text{ V})/(5.0 \Omega)$$

For computing the correct answer

1 point

$$I = 0.090 \text{ A}$$

For indicating a clockwise direction for the current

1 point

## (c) 3 points

For relating the energy dissipated to the power in the resistor

1 point

$$E = \int P dt \quad \text{or} \quad E = Pt$$

For an expression for electric power

1 point

$$P = I^2 R \quad \text{or} \quad \frac{V^2}{R} \quad \text{or} \quad IV$$

Example using  $P = I^2 R$ :

$$E = I^2 Rt$$

$$E = (0.090 \text{ A})^2 (5.0 \Omega)(15 \text{ s})$$

For computing the correct answer

1 point

$$E = 0.61 \text{ J}$$

**1999 Physics C Solutions****Distribution  
of Points**

E &amp; M 2 (continued)

(d) 3 points

For stating that the brightness of the bulb will be less

1 point

For indicating that the reduction in brightness is due to a decrease in current or a decrease in the emf

1 point

For indicating that the decrease in current or emf, or the reduction in brightness, is due to a decrease in the area of the loop or a decrease in the changing flux

1 point

For using correct units with three numerical answers

1 point

E &amp; M 3 (15 points)

(a) 3 points

The charge on any section of the ring is equidistant from a point on the  $x$ -axis,  
so one can write an equation in terms of the single distance  $r$

For a correct expression of the potential

$$= \frac{1}{4\pi\epsilon_0} \frac{dq}{r} \quad \text{or} \quad \frac{1}{4\pi\epsilon_0} \frac{Q}{r}$$

1 point

For a correct expression for the distance of the charge from location  $x$

$$r = \sqrt{x^2 + R^2}$$

1 point

For the correct answer

$$= \frac{1}{4\pi\epsilon_0} \frac{Q}{\sqrt{x^2 + R^2}}$$

1 point

*Alternate solution*

*Alternate points*

For correctly expressing the potential as an integral of the electric field

1 point

$$dV = -\int E \, dr$$

For a correct expression for the field

1 point

$$dV = -\int \frac{1}{4\pi\epsilon_0} \frac{Qx}{(x^2 + R^2)^{3/2}} dx$$

For correctly integrating to get the final answer

1 point

$$= \frac{1}{4\pi\epsilon_0} \frac{Q}{\sqrt{x^2 + R^2}}$$

(b)

i. 3 points

$$E = -\frac{dV}{dr}$$

For using the above relationship

1 point

For taking the derivative with respect to  $x$

1 point

For using the expression for  $V$  obtained in part (a)

1 point

$$E_x = -\frac{d}{dx} \left( \frac{1}{4\pi\epsilon_0} \frac{Q}{\sqrt{x^2 + R^2}} \right)$$

$$E_x = \frac{1}{4\pi\epsilon_0} \frac{Qx}{(x^2 + R^2)^{3/2}}$$

## 1999 Physics C Solutions

Distribution  
of points

E &amp; M 3 (continued)

(b) (continued)

i. (continued)

*Alternate solution**Alternate points*

Calculating the field by integration:

$$E = \int dE_x = \int dE \cos \theta, \text{ where } \theta \text{ is the angle between the } x\text{-axis and}$$

the distance vector  $\mathbf{r}$

For using the horizontal component of the field

1 point

For using a correct expression of Coulomb's law

1 point

$$E = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}$$

For indicating that the integral is over the charge

1 point

$$E_x = \int \frac{1}{4\pi\epsilon_0} \frac{dq}{r^2} \cos \theta$$

Substituting  $\cos \theta = x/r$  and  $r = \sqrt{x^2 + R^2}$ 

$$E_x = \frac{1}{4\pi\epsilon_0} \frac{Q}{(x^2 + R^2)^{3/2}} \int_0^Q dq$$

$$E_x = \frac{1}{4\pi\epsilon_0} \frac{Qx}{(x^2 + R^2)^{3/2}}$$

ii. 1 point

For any indication that the  $y$ - and  $z$ -components are zero or cancel

1 point

## E &amp; M 3 (continued)

(c)

i. 2 points

For taking the derivative of  $E$  with respect to  $x$  and setting it equal to zero

1 point

$$\frac{dE_x}{dx} = \frac{d}{dx} \left( \frac{1}{4\pi\epsilon_0} \frac{Qx}{(x^2 + R^2)^{3/2}} \right) = 0$$

$$\frac{Q}{4\pi\epsilon_0} \left( \frac{1}{(x^2 + R^2)^{3/2}} + \left(-\frac{3}{2}\right) \frac{2x^2}{(x^2 + R^2)^{5/2}} \right) = 0$$

$$\frac{1}{(x^2 + R^2)^{3/2}} - \frac{3x^2}{(x^2 + R^2)^{5/2}} = 0$$

$$\frac{1}{(x^2 + R^2)^{3/2}} = \frac{3x^2}{(x^2 + R^2)^{5/2}}$$

$$1 = \frac{3x^2}{x^2 + R^2}$$

$$3x^2 = x^2 + R^2$$

$$x = \pm \frac{R}{\sqrt{2}} \quad \text{and the maximum occurs at the positive value of } x$$

For the correct answer

1 point

$$x = + \frac{R}{\sqrt{2}}$$

ii. 1 point

For substituting the answer from part (c)i into the given expression for the electric field


1 point

$$E_{x \text{ max}} = \frac{1}{4\pi\epsilon_0} \frac{Q(R/\sqrt{2})}{\left((R/\sqrt{2})^2 + R^2\right)^{3/2}}$$

$$E_{x \text{ max}} = \frac{1}{4\pi\epsilon_0} \frac{2Q}{3\sqrt{3}R^2}$$

E &amp; M 3 (continued)

(d) 3 points



- |   |         |
|---|---------|
| For a curve in the first quadrant displaying a single positive maximum        | 1 point |
| For a curve passing through the origin  | 1 point |
| For the negative reflection of the first quadrant curve in the third quadrant | 1 point |

(e) 2 points

- |  |          |
|--|----------|
| For any statement that describes the subsequent motion as oscillating, periodic etc. | 2 points |
|--|----------|

One point was awarded for a statement that only described the electron as moving toward the ring or along the  $x$ -axis.