



## AP Physics C: Electricity and Magnetism 2001 Scoring Guidelines

**The materials included in these files are intended for non-commercial use by AP teachers for course and exam preparation; permission for any other use must be sought from the Advanced Placement Program. Teachers may reproduce them, in whole or in part, in limited quantities, for face-to-face teaching purposes but may not mass distribute the materials, electronically or otherwise. These materials and any copies made of them may not be resold, and the copyright notices must be retained as they appear here. This permission does not apply to any third-party copyrights contained herein.**

These materials were produced by Educational Testing Service (ETS), which develops and administers the examinations of the Advanced Placement Program for the College Board. The College Board and Educational Testing Service (ETS) are dedicated to the principle of equal opportunity, and their programs, services, and employment policies are guided by that principle.

The College Board is a national nonprofit membership association dedicated to preparing, inspiring, and connecting students to college and opportunity. Founded in 1900, the association is composed of more than 3,900 schools, colleges, universities, and other educational organizations. Each year, the College Board serves over three million students and their parents, 22,000 high schools, and 3,500 colleges, through major programs and services in college admission, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT<sup>®</sup>, the PSAT/NMSQT<sup>™</sup>, the Advanced Placement Program<sup>®</sup> (AP<sup>®</sup>), and Pacesetter<sup>®</sup>. The College Board is committed to the principles of equity and excellence, and that commitment is embodied in all of its programs, services, activities, and concerns.

Copyright © 2001 by College Entrance Examination Board. All rights reserved. College Board, Advanced Placement Program, AP, and the acorn logo are registered trademarks of the College Entrance Examination Board.

# AP<sup>®</sup> PHYSICS C: ELECTRICITY AND MAGNETISM

## 2001 SCORING GUIDELINES

### General Notes about 2001 AP Physics Solutions

1. The solutions contain the most common method(s) of solving the free-response questions, and the allocation of points for these solutions. Other methods of solution also receive appropriate credit for correct work.
2. Generally, double penalty for errors is avoided. For example, if an incorrect answer to part (a) is correctly substituted into an otherwise correct solution to part (b), full credit will usually be awarded.
3. An exception to this may be cases when the numerical answer to a later part should be easily recognized as wrong, e.g., a speed faster than the speed of light in vacuum.
4. Implicit statements of concepts normally receive credit. For example, if use of the equation expressing a particular concept is worth one point, and a solution contains the application of the equation to the problem but does not separately list the basic equation, the point is still awarded.

**AP<sup>®</sup> PHYSICS C: ELECTRICITY AND MAGNETISM  
2001 SCORING GUIDELINES**

**Question 1**

**15 points total**

1. (a) **4 points**

**Distribution  
of Points**

For a correct formula for determining the electric field

**1 point**

$$\mathbf{E} = \sum \frac{kQ}{r^2} \hat{\mathbf{r}} \quad \text{or} \quad \mathbf{E} = \frac{kQ}{r^2} \hat{\mathbf{r}}$$

Summing the contributions to the field from the four charges, letting fields directed upward be positive and fields directed downward be negative (full credit also given for using opposite convention as long as answers were consistent):

$$E = - \frac{(9 \times 10^9)(30)}{(3 \times 10^3)^2} + \frac{(9 \times 10^9)(30)}{(2 \times 10^3)^2} + \frac{(9 \times 10^9)(30)}{(2 \times 10^3)^2} - \frac{(9 \times 10^9)(30)}{(3 \times 10^3)^2}$$

For correct substitutions shown in the above equation

**1 point**

$$E = -30,000 \text{ N/C} + 67,500 \text{ N/C} + 67,500 \text{ N/C} - 30,000 \text{ N/C}$$

$$E = 75,000 \text{ N/C, directed upward}$$

**1 point**

For correct magnitude ( $E = \frac{8 \times 10^{-6}}{4\pi\epsilon_0}$  or  $k(8 \times 10^{-6})$  also accepted)

For correct direction, either stated or shown by an upward directed arrow

**1 point**

Notes: If wrong signs were used in the substitution, the point for correct magnitude was not awarded. If calculation was done using only the two real charges, a maximum of **3 points** was awarded as follows: **1 point** for the formula, **1 point** for the calculation, **1 point** for direction.

1. (b) i. **1 point**

For correctly indicating direction, such as by an upward directed arrow at  $P_2$ ,

**1 point**

1. (b) ii. **2 points**

For correctly checking the space in front of “Less”

**1 point**

For correct justification, such as “ $P_2$  is farther from all the charges than  $P_1$ , so the net field is less.” If student chose to work out the actual magnitude of the field at  $P_2$  (which is about 45,000 N/C), the justification point was awarded for any calculated numerical value less than 75,000 N/C.

**1 point**

Note: **No points** were awarded for (b) ii. if the wrong space was checked.

**AP<sup>®</sup> PHYSICS C: ELECTRICITY AND MAGNETISM  
2001 SCORING GUIDELINES**

**Question 1 (cont.)**

1. (c) i. **2 points**

**Distribution  
of Points**

The potential at  $P_1$  is 0, which can be determined without calculation from symmetry considerations.

For stating  $V = 0$  or for just 0

**2 points**

*Alternate Solution*

*Alternate  
points*

For a correct formula for determining the potential

*1 point*

$$V = \sum \frac{kQ}{r}$$

$$V = k \left( \frac{30}{3000} - \frac{30}{2000} + \frac{30}{2000} - \frac{30}{3000} \right) = 0$$

For the correct answer

*1 point*

1. (c) ii. **2 points**

The potential at  $P_2$  is 0, which can also be determined without calculation from symmetry considerations.

For stating  $V = 0$  or for just 0

**2 points**

*Alternate Solution*

*Alternate  
points*

For a correct formula for determining the potential

*1 point*

$$V = \sum \frac{kQ}{r}$$

$$V = k \left( \frac{30}{\sqrt{10} \times 10^3} - \frac{30}{\sqrt{5} \times 10^3} + \frac{30}{\sqrt{5} \times 10^3} - \frac{30}{\sqrt{10} \times 10^3} \right) = 0$$

For the correct answer

*1 point*

1. (d) **2 points**

For a correct formula for determining the potential

**1 point**

$$V = \sum \frac{kQ}{r}$$

For correct substitution of values (no credit lost for failing to convert kilometers to meters)

**1 point**

$$V_p = k \left( \frac{30}{(3-1) \times 10^3} - \frac{30}{(2-1) \times 10^3} + \frac{30}{(2+1) \times 10^3} - \frac{30}{(3+1) \times 10^3} \right)$$

$$V_p = -1.12 \times 10^8 \text{ V} \approx -10^8 \text{ V}$$

**AP<sup>®</sup> PHYSICS C: ELECTRICITY AND MAGNETISM  
2001 SCORING GUIDELINES**

**Question 1 (cont.)**

1. (e) **2 points**

**Distribution  
of Points**

For a correct formula for the potential energy (must include a summation sign)

**1 point**

$$U = k \sum_{i,j} \frac{q_i q_j}{r_{ij}} \quad \text{or} \quad U = \sum qV$$

For correct substitution without regard to sign error (no credit lost for failing to convert kilometers to meters)

**1 point**

$$U = (9 \times 10^9) \left[ \frac{30(-30)}{10^3} + \frac{30(30)}{5 \times 10^3} + \frac{30(-30)}{6 \times 10^3} + \frac{-30(30)}{4 \times 10^3} + \frac{-30(-30)}{5 \times 10^3} + \frac{30(-30)}{10^3} \right] \text{ J}$$

$$U = -1.6 \times 10^{10} \text{ J}$$

**AP<sup>®</sup> PHYSICS C: ELECTRICITY AND MAGNETISM  
2001 SCORING GUIDELINES**

**Question 2**

**15 points total**

2. (a) **4 points**

**Distribution  
of Points**

There were three methods generally used to solve this problem.

Method 1.

For a correct method based on determining the time constant using values from the graph **2 points**

$$\tau = RC \approx 60 \text{ min} = 3600 \text{ s}$$

For correct substitution of values with proper units **1 point**

$$R = \frac{\tau}{C} = \frac{3600 \text{ s}}{8.0 \times 10^{-6} \text{ F}}$$

For answer consistent with values used **1 point**

$$R = 4.5 \times 10^8 \Omega$$

Method 2.

$$V = V_0 e^{-t/RC}$$

For using the above equation with given value of  $C$  and values for  $V$ ,  $V_0$ , and  $t$  from the graph with  $t$  correlating with  $V$  **2 points**

Example:  $V_0 = 10 \text{ V}$ , and  $V = 2 \text{ V}$  at  $t = 100 \text{ min}$

For correct substitution of values **1 point**

$$2 = 10 e^{(-6000 \text{ s})/R(8 \times 10^{-6} \text{ F})}$$

$$\ln(2/10) = (6000)R(8 \times 10^{-6})$$

For answer consistent with values used **1 point**

$$R = 4.7 \times 10^8 \Omega$$

Method 3.

Find a correct relationship that depends on the slope of the graph:

Example:  $R = \frac{V}{i} = \frac{V}{dQ/dt}$

But  $dQ = CdV$

So  $R = \frac{V}{C(dV/dt)}$

For estimating  $dV/dt$  by computing  $\Delta V/\Delta t$  for a particular value of  $V$  **2 points**

For substituting values in equation above **1 point**

For answer consistent with values used **1 point**

Note: The value used for  $V$  must be that at the point where the slope is taken and clearly indicated.

**AP<sup>®</sup> PHYSICS C: ELECTRICITY AND MAGNETISM  
2001 SCORING GUIDELINES**

**Question 2 (cont.)**

2. (b) **3 points**

**Distribution  
of Points**

For a correct equation for the capacitance

**1 point**

$$C = \frac{\kappa\epsilon_0 A}{d}$$

For correct algebraic solution for area  $A$  and substitution of variables

**1 point**

$$A = \frac{Cd}{\kappa\epsilon_0} = \frac{(8.0 \times 10^{-6} \text{ F})(1.0 \times 10^{-4} \text{ m})}{5.6(8.85 \times 10^{-12} \text{ C}^2/\text{N} \cdot \text{m})}$$

For the correct answer

**1 point**

$$A = 16 \text{ m}^2$$

2. (c) **3 points**

For a correct equation for the resistance

**1 point**

$$R = \frac{\rho L}{A}$$

For correct algebraic solution for resistivity  $\rho$  and substitution of variables

**1 point**

$$\rho = \frac{RA}{L} = \frac{(4.5 \times 10^8 \text{ } \Omega)(16 \text{ m}^2)}{1.0 \times 10^{-4} \text{ m}}$$

For answer consistent with values used

**1 point**

$$\rho = 7.2 \times 10^{13} \text{ } \Omega \cdot \text{m}$$

**AP<sup>®</sup> PHYSICS C: ELECTRICITY AND MAGNETISM  
2001 SCORING GUIDELINES**

**Question 2 (cont.)**

2. (d) **4 points**

**Distribution  
of Points**

There were four general methods for solving this problem. Solutions were scored as follows:

For using a correct method

**2 points**

For substituting appropriate values

**1 point**

For answer consistent with values obtained in earlier parts

**1 point**

Method 1:

$\Delta Q = C\Delta V$ , and substitute values obtained from graph

Example:  $Q_i = (8.0 \times 10^{-6} \text{ F})(10 \text{ V})$  and  $Q_f = (8.0 \times 10^{-6} \text{ F})(2 \text{ V})$

So  $\Delta Q = 6.4 \times 10^{-5} \text{ C}$  or  $64 \mu\text{C}$

Method 2:

$$Q = \int_0^{6000} I dt = \frac{V_0}{R} \int_0^{6000} e^{-t/RC} dt = \frac{10 \text{ V}}{4.5 \times 10^8 \Omega} \int_0^{6000} e^{-t/3600} dt$$

$$Q = (2.22 \times 10^{-8})(-3600)e^{-t/3600} \Big|_0^{6000 \text{ s}}$$

$$Q = 6.5 \times 10^{-5} \text{ C or } 65 \mu\text{C}$$

Method 3:

$$Q = CV_0(1 - e^{-t/RC})$$

$$Q = (8.0 \times 10^{-6} \text{ F})(10 \text{ V}) \left( 1 - e^{-6000 \text{ s} / (4.5 \times 10^8 \Omega)(8.0 \times 10^{-6} \text{ F})} \right)$$

$$Q = 6.5 \times 10^{-5} \text{ C or } 65 \mu\text{C}$$

Method 4:

Determine the area under the curve of  $I$  vs.  $t$ , which is the  $V$  vs.  $t$  graph shown with  $I = V_0/R$ .

Each block has area equal to  $Vt/R = 1.3 \times 10^{-6} \text{ C}$ .

Estimating 47 blocks under the curve gives

$$Q = 6.3 \times 10^{-5} \text{ C or } 63 \mu\text{C}$$

**Unit point:** For correct units given on answers for **three** of the **four** parts

**1 point**

**AP<sup>®</sup> PHYSICS C: ELECTRICITY AND MAGNETISM  
2001 SCORING GUIDELINES**

**Question 3**

**15 points total**

3. (a) **2 points**

**Distribution  
of Points**

Using Ohm's law:

$$V = IR$$

For correct equation for  $I$

**1 point**

$$I = \frac{\mathcal{E}}{R}$$

For correctly indicating the current direction on the diagram or in the answer space, such as by stating that it is clockwise, or to the left, or by showing an arrow pointing left

**1 point**

3. (b) **4 points**

For indicating on the diagram or in the answer space a direction opposite to the answer in part (a). If part (a) does not contain a direction, then for an indication that the direction is to the right or by showing an arrow pointing right.

**1 point**

For a complete justification

**3 points**

**Full credit** awarded for an answer that indicated the right-hand rule to obtain the magnetic field directed out of the page at the rod, and then used the cross product to obtain that the force on the rod is up

**2 points** partial credit awarded for an answer that just stated the rule that antiparallel currents repel or that just stated  $I \ell \times \mathbf{B}$  and the right-hand rule

**1 point** partial credit awarded for an answer that just stated the right-hand rule or  $I \ell \times \mathbf{B}$  or some fragment with some correct element

3. (c) **4 points**

For indicating that the gravitational force will be equal to  $I \ell \times \mathbf{B}$

**1 point**

$$F = I \ell \times \mathbf{B} = mg$$

For giving the correct equation for the magnetic field

**1 point**

$$B = \frac{\mu_0 i}{2\pi r}$$

For correctly substituting in the first equation above the values for  $B$  and for  $I$  from part (a)

**1 point**

$$\frac{\mu_0 I_c \ell \mathcal{E}}{2\pi r R} = mg$$

For the correct answer

**1 point**

$$I_c = \frac{2\pi mgrR}{\mu_0 \ell \mathcal{E}}$$

**AP<sup>®</sup> PHYSICS C: ELECTRICITY AND MAGNETISM  
2001 SCORING GUIDELINES**

**Question 3 (cont.)**

3. (d) **5 points**

**Distribution  
of Points**

For the correct expression for  $\phi$

**1 point**

$\phi = \int \mathbf{B} \cdot d\mathbf{A}$ , where  $B = \frac{\mu_0 I_c}{2\pi x}$ , and  $x$  is the vertical distance from the cable

Letting  $dA = \ell dx$  and substituting the values for  $B$  above and for  $I_c$  from part (c):

$$\phi = \int_r^{r+d} \frac{\mu_0 2\pi mgrR\ell}{2\pi\mu_0\ell\mathcal{E}} \frac{dx}{x}$$

For correct limits of integration

**1 point**

For correct substitution of the values consistent with previous answers

**1 point**

For correct integration

**1 point**

$$\phi = \frac{mgrR}{\mathcal{E}} \ln x \Big|_r^{r+d}$$

For the correct answer

**1 point**

$$\phi = \frac{mgrR}{\mathcal{E}} \ln\left(\frac{r+d}{r}\right)$$