



## AP Physics C: Electricity and Magnetism 2000 Student Samples

**The materials included in these files are intended for non-commercial use by AP teachers for course and exam preparation; permission for any other use must be sought from the Advanced Placement Program. Teachers may reproduce them, in whole or in part, in limited quantities, for face-to-face teaching purposes but may not mass distribute the materials, electronically or otherwise. These materials and any copies made of them may not be resold, and the copyright notices must be retained as they appear here. This permission does not apply to any third-party copyrights contained herein.**

These materials were produced by Educational Testing Service (ETS), which develops and administers the examinations of the Advanced Placement Program for the College Board. The College Board and Educational Testing Service (ETS) are dedicated to the principle of equal opportunity, and their programs, services, and employment policies are guided by that principle.

The College Board is a national nonprofit membership association dedicated to preparing, inspiring, and connecting students to college and opportunity. Founded in 1900, the association is composed of more than 3,900 schools, colleges, universities, and other educational organizations. Each year, the College Board serves over three million students and their parents, 22,000 high schools, and 3,500 colleges, through major programs and services in college admission, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT<sup>®</sup>, the PSAT/NMSQT<sup>™</sup>, the Advanced Placement Program<sup>®</sup> (AP<sup>®</sup>), and Pacesetter<sup>®</sup>. The College Board is committed to the principles of equity and excellence, and that commitment is embodied in all of its programs, services, activities, and concerns.

Copyright © 2001 by College Entrance Examination Board. All rights reserved. College Board, Advanced Placement Program, AP, and the acorn logo are registered trademarks of the College Entrance Examination Board.

E & M 2.

Three particles,  $A$ ,  $B$ , and  $C$ , have equal positive charges  $Q$  and are held in place at the vertices of an equilateral triangle with sides of length  $\ell$ , as shown in the figures below. The dotted lines represent the bisectors for each side. The base of the triangle lies on the  $x$ -axis, and the altitude of the triangle lies on the  $y$ -axis.




Figure 1




Figure 2

- (a)
- Point  $P_1$ , the intersection of the three bisectors, locates the geometric center of the triangle and is one point where the electric field is zero. On Figure 1 above, draw the electric field vectors  $E_A$ ,  $E_B$ , and  $E_C$  at  $P_1$  due to each of the three charges. Be sure your arrows are drawn to reflect the relative magnitude of the fields.
  - Another point where the electric field is zero is point  $P_2$  at  $(0, y_2)$ . On Figure 2 above, draw electric field vectors  $E_A$ ,  $E_B$ , and  $E_C$  at  $P_2$  due to each of the three point charges. Indicate below whether the magnitude of each of these vectors is greater than, less than, or the same as for point  $P_1$ .

	Greater than at $P_1$	Less than at $P_1$	The same as at $P_1$
$E_A$	No	Yes	No
$E_B$	Yes	No	No
$E_C$	Yes	No	No

GO ON TO THE NEXT PAGE.



# EEEEEEEEEEEEEEEEEE \ 2

(b) Explain why the x-component of the total electric field is zero at any point on the y-axis.

Because on the y-axis, the x-component of the field from A will always be 0 since it is directly above. The x-components from B & C will cancel each other out since they are equidistant from whatever point on the y-axis is chosen and they are on opposite sides. Therefore, there is no x-component of the total electric field on any point on the y-axis.

(c) Write a general expression for the electric potential  $V$  at any point on the y-axis inside the triangle in terms of  $Q$ ,  $\ell$ , and  $y$ .

$$V = \frac{1}{4\pi\epsilon_0} \left( \frac{Q}{\left(\frac{\ell}{2} - y\right)} + \frac{Q}{\sqrt{y^2 + \frac{\ell^2}{4}}} + \frac{Q}{\sqrt{y^2 + \frac{\ell^2}{4}}} \right)$$



$$\left(\frac{\ell}{2}\right)^2 - \left(\frac{\ell}{2}\right)^2 = \frac{\ell^2}{4}$$

$$\sqrt{y^2 + \frac{\ell^2}{4}}$$

(d) Describe how the answer to part (c) could be used to determine the y-coordinates of points  $P_1$  and  $P_2$  at which the electric field is zero. (You do not need to actually determine these coordinates.)

When the electric field is zero, the opposite of the derivative of the voltage is equal to zero ( $E = -\frac{dV}{dy}$ ). Find the derivative of  $V$ , set it equal to zero, and solve for  $y$  to get the y-coordinates at which the electric field is zero.

GO ON TO THE NEXT PAGE.

E & M 2.

Three particles,  $A$ ,  $B$ , and  $C$ , have equal positive charges  $Q$  and are held in place at the vertices of an equilateral triangle with sides of length  $\ell$ , as shown in the figures below. The dotted lines represent the bisectors for each side. The base of the triangle lies on the  $x$ -axis, and the altitude of the triangle lies on the  $y$ -axis.




Figure 1




Figure 2

(a)

- i. Point  $P_1$ , the intersection of the three bisectors, locates the geometric center of the triangle and is one point where the electric field is zero. On Figure 1 above, draw the electric field vectors  $\mathbf{E}_A$ ,  $\mathbf{E}_B$ , and  $\mathbf{E}_C$  at  $P_1$  due to each of the three charges. Be sure your arrows are drawn to reflect the relative magnitude of the fields.
- ii. Another point where the electric field is zero is point  $P_2$  at  $(0, y_2)$ . On Figure 2 above, draw electric field vectors  $\mathbf{E}_A$ ,  $\mathbf{E}_B$ , and  $\mathbf{E}_C$  at  $P_2$  due to each of the three point charges. Indicate below whether the magnitude of each of these vectors is greater than, less than, or the same as for point  $P_1$ .

	Greater than at $P_1$	Less than at $P_1$	The same as at $P_1$
$E_A$		✓	
$E_B$	✓		
$E_C$	✓		

GO ON TO THE NEXT PAGE.

(b) Explain why the x-component of the total electric field is zero at any point on the y-axis.

The x-components of the  $E_B$  &  $E_C$  vectors will always be equal & opposite, and the x-component of the  $E_A$  vector is zero, so the vector sum of the 3 will always be zero.

(c) Write a general expression for the electric potential  $V$  at any point on the y-axis inside the triangle in terms of  $Q$ ,  $l$ , and  $y$ .

distance from A =  $l\frac{\sqrt{3}}{2} - y$

distance from B = distance from C =  $\sqrt{\frac{l^2}{4} + y^2}$

voltage =  $\frac{1}{4\pi\epsilon_0} \frac{Q}{R}$ , and voltages are additive, so

$$V = \frac{1}{4\pi\epsilon_0} Q \left[ \frac{1}{l\frac{\sqrt{3}}{2} - y} + \frac{2}{\sqrt{\frac{l^2}{4} + y^2}} \right]$$

(d) Describe how the answer to part (c) could be used to determine the y-coordinates of points  $P_1$  and  $P_2$  at which the electric field is zero. (You do not need to actually determine these coordinates.)

~~the~~  $\vec{E} = -\nabla V$ ; on the x-axis that means

$$\vec{E} = -dV/dx. \quad \text{if } |\vec{E}| = 0,$$

$$|-dV/dx| = 0, \text{ so}$$

we can take  $dV/dx$ , set it = 0, and solve.



