



AP Physics C: Mechanics 2000 Student Samples

The materials included in these files are intended for non-commercial use by AP teachers for course and exam preparation; permission for any other use must be sought from the Advanced Placement Program. Teachers may reproduce them, in whole or in part, in limited quantities, for face-to-face teaching purposes but may not mass distribute the materials, electronically or otherwise. These materials and any copies made of them may not be resold, and the copyright notices must be retained as they appear here. This permission does not apply to any third-party copyrights contained herein.

These materials were produced by Educational Testing Service (ETS), which develops and administers the examinations of the Advanced Placement Program for the College Board. The College Board and Educational Testing Service (ETS) are dedicated to the principle of equal opportunity, and their programs, services, and employment policies are guided by that principle.

The College Board is a national nonprofit membership association dedicated to preparing, inspiring, and connecting students to college and opportunity. Founded in 1900, the association is composed of more than 3,900 schools, colleges, universities, and other educational organizations. Each year, the College Board serves over three million students and their parents, 22,000 high schools, and 3,500 colleges, through major programs and services in college admission, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT[®], the PSAT/NMSQT[™], the Advanced Placement Program[®] (AP[®]), and Pacesetter[®]. The College Board is committed to the principles of equity and excellence, and that commitment is embodied in all of its programs, services, activities, and concerns.

Copyright © 2001 by College Entrance Examination Board. All rights reserved. College Board, Advanced Placement Program, AP, and the acorn logo are registered trademarks of the College Entrance Examination Board.

PHYSICS C
Section II, MECHANICS
Time—45 minutes
3 Questions

Directions: Answer all three questions. The suggested time is about 15 minutes for answering each of the questions, which are worth 15 points each. The parts within a question may not have equal weight. Show all your work in this booklet in the spaces provided after each part, NOT in the green insert.

Mech 1.

You are conducting an experiment to measure the acceleration due to gravity g_0 at an unknown location. In the measurement apparatus, a simple pendulum swings past a photogate located at the pendulum's lowest point, which records the time t_{10} for the pendulum to undergo 10 full oscillations. The pendulum consists of a sphere of mass m at the end of a string and has a length ℓ . There are four versions of this apparatus, each with a different length. All four are at the unknown location, and the data shown below are sent to you during the experiment.

ℓ (cm)	t_{10} (s)	T (s)	T^2 (s ²)
12	7.62	.762 s	.581
18	8.89	.889 s	.790
21	10.09	1.009 s	1.018
32	12.08	1.208 s	1.459

- (a) For each pendulum, calculate the period T and the square of the period. Use a reasonable number of significant figures. Enter these results in the table above.

$$T = \frac{t_{10}}{10} = \frac{7.62 \text{ s}}{10} = .762 \text{ s}$$


$$\frac{8.89}{10} = .889 \text{ s}$$

etc...

$$T^2 = (.762 \text{ s})^2 = .581 \text{ s}^2$$

etc...

(b) On the axes below, plot the square of the period versus the length of the pendulum. Draw a best-fit straight line for this data.



(c) Assuming that each pendulum undergoes small amplitude oscillations, from your fit determine the experimental value g_{exp} of the acceleration due to gravity at this unknown location. Justify your answer.

$$T = 2\pi \sqrt{\frac{l}{g}}$$

$$T^2 = \frac{4\pi^2 l}{g} \quad g = \frac{4\pi^2 l}{T^2} = \frac{4\pi^2}{m} \quad (1)$$

From graph: $m = \frac{\Delta(T^2)}{\Delta l} = \frac{1.48 \text{ s}^2 - 0 \text{ s}^2}{32 \text{ cm} - 0} = 4.63; \frac{\text{s}^2}{\text{m}}$

Substitute m into (1):


$$g = \frac{4\pi^2}{m} = \frac{4\pi^2}{4.63 \frac{\text{s}^2}{\text{m}}} = \boxed{8.53; \frac{\text{m}}{\text{s}^2}}$$

(d) If the measurement apparatus allows a determination of g_{exp} that is accurate to within 4%, is your experimental value in agreement with the value 9.80 m/s^2 ? Justify your answer.

$$\begin{aligned} \% \text{ diff} &= \frac{\text{exp} - \text{act}}{\text{act}} (100\%) \\ &= \frac{9.53 - 9.80}{9.80} (100\%) \\ &= -12.9\% \end{aligned}$$

No, the value is not in agreement, it is almost 13% different.

(e) Someone informs you that the experimental apparatus is in fact near Earth's surface, but that the experiment has been conducted inside an elevator with a constant acceleration a . Assuming that your experimental value g_{exp} is exact, determine the magnitude and direction of the elevator's acceleration.



If elevator was in freefall, g felt by occupant is zero. If elevator ascends, g felt by occupants is greater than 9.80

Because $g_{\text{exp}} = 9.53 \frac{\text{m}}{\text{s}^2}$, $a_{\text{elev}} = 9.8 \frac{\text{m}}{\text{s}^2} - 9.53 \frac{\text{m}}{\text{s}^2}$

$$0 < a_{\text{elev}} < 9.8 \implies \boxed{= 1.27 \frac{\text{m}}{\text{s}^2} \downarrow}$$

PHYSICS C
Section II, MECHANICS
 Time—45 minutes
 3 Questions

Directions: Answer all three questions. The suggested time is about 15 minutes for answering each of the questions, which are worth 15 points each. The parts within a question may not have equal weight. Show all your work in this booklet in the spaces provided after each part, NOT in the green insert.


Mech 1.

You are conducting an experiment to measure the acceleration due to gravity g_0 at an unknown location. In the measurement apparatus, a simple pendulum swings past a photogate located at the pendulum's lowest point, which records the time t_{10} for the pendulum to undergo 10 full oscillations. The pendulum consists of a sphere of mass m at the end of a string and has a length ℓ . There are four versions of this apparatus, each with a different length. All four are at the unknown location, and the data shown below are sent to you during the experiment.

ℓ (cm)	t_{10} (s)	T (s)	T^2 (s ²)
12	7.62	.762	.581
18	8.89	.889	.790
21	10.09	1.01	1.02
32	12.08	1.21	1.46

- (a) For each pendulum, calculate the period T and the square of the period. Use a reasonable number of significant figures. Enter these results in the table above.

(b) On the axes below, plot the square of the period versus the length of the pendulum. Draw a best-fit straight line for this data.



(c) Assuming that each pendulum undergoes small amplitude oscillations, from your fit determine the experimental value g_{exp} of the acceleration due to gravity at this unknown location. Justify your answer.

$$T_p = 2\pi\sqrt{\frac{l}{g}}, \text{ so } T_p^2 = 4\pi^2 \cdot \frac{l}{g}$$

$$g = \frac{4\pi^2 l}{T^2} \quad \frac{l}{T^2} \approx 22 \text{ (marked "x" above)}$$

$$g = 22 \cdot 4\pi^2 = 88\pi^2 = 869 \frac{\text{cm}}{\text{s}^2}$$

$$\left(= 8.69 \frac{\text{m}}{\text{s}^2} \right)$$


GO ON TO THE NEXT PAGE.

- (d) If the measurement apparatus allows a determination of g_u that is accurate to within 4%, is your experimental value in agreement with the value 9.80 m/s^2 ? Justify your answer.

$$\frac{9.80 - 8.69}{9.80} = 11.3\% \text{ error}$$

not w/in 4%, so g_{exp} is
not in agreement

- (e) Someone informs you that the experimental apparatus is in fact near Earth's surface, but that the experiment has been conducted inside an elevator with a constant acceleration a . Assuming that your experimental value g_{exp} is exact, determine the magnitude and direction of the elevator's acceleration.



PHYSICS C
Section II, MECHANICS
Time—45 minutes
3 Questions

Directions: Answer all three questions. The suggested time is about 15 minutes for answering each of the questions, which are worth 15 points each. The parts within a question may not have equal weight. Show all your work in this booklet in the spaces provided after each part, NOT in the green insert.

Mech 1.

You are conducting an experiment to measure the acceleration due to gravity g_0 at an unknown location. In the measurement apparatus, a simple pendulum swings past a photogate located at the pendulum's lowest point, which records the time t_{10} for the pendulum to undergo 10 full oscillations. The pendulum consists of a sphere of mass m at the end of a string and has a length ℓ . There are four versions of this apparatus, each with a different length. All four are at the unknown location, and the data shown below are sent to you during the experiment.


ℓ (cm)	t_{10} (s)	T (s)	T^2 (s ²)
12	7.62	0.762	0.580644
18	8.89	0.889	0.790321
21	10.09	1.009	1.018081
32	12.08	1.208	1.459264

- (a) For each pendulum, calculate the period T and the square of the period. Use a reasonable number of significant figures. Enter these results in the table above.

$$T_p = 2\pi \sqrt{\frac{\ell}{g}}$$

$$\frac{t_{10}}{10} = T_p$$

(b) On the axes below, plot the square of the period versus the length of the pendulum. Draw a best-fit straight line for this data.



(c) Assuming that each pendulum undergoes small amplitude oscillations, from your fit determine the experimental value g_{exp} of the acceleration due to gravity at this unknown location. Justify your answer.

$$T_p = 2\pi \sqrt{\frac{l}{g}}$$

$$T_p^2 = 2^2 \pi^2 \frac{l}{g}$$

$$0.580644 = 4\pi^2 \times \frac{12}{g}$$

$$g = 8.16 \text{ m/s}^2$$

- (d) If the measurement apparatus allows a determination of g_{ex} that is accurate to within 4%, is your experimental value in agreement with the value 9.80 m/s^2 ? Justify your answer.

Nope. I calculated g to be 8.16 m/s^2

$$\frac{9.8 - 8.16}{9.8} \times 100 = 16\% \text{ error}$$

- (c) Someone informs you that the experimental apparatus is in fact near Earth's surface, but that the experiment has been conducted inside an elevator with a constant acceleration a . Assuming that your experimental value g_{exp} is exact, determine the magnitude and direction of the elevator's acceleration.