



AP[®] Physics C: Electricity & Magnetism 1999 Sample Student Responses

The materials included in these files are intended for non-commercial use by AP teachers for course and exam preparation; permission for any other use must be sought from the Advanced Placement Program. Teachers may reproduce them, in whole or in part, in limited quantities, for face-to-face teaching purposes but may not mass distribute the materials, electronically or otherwise. These materials and any copies made of them may not be resold, and the copyright notices must be retained as they appear here. This permission does not apply to any third-party copyrights contained herein.

These materials were produced by Educational Testing Service (ETS), which develops and administers the examinations of the Advanced Placement Program for the College Board. The College Board and Educational Testing Service (ETS) are dedicated to the principle of equal opportunity, and their programs, services, and employment policies are guided by that principle.

The College Board is a national nonprofit membership association dedicated to preparing, inspiring, and connecting students to college and opportunity. Founded in 1900, the association is composed of more than 3,900 schools, colleges, universities, and other educational organizations. Each year, the College Board serves over three million students and their parents, 22,000 high schools, and 3,500 colleges, through major programs and services in college admission, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT[®], the PSAT/NMSQT[™], the Advanced Placement Program[®] (AP[®]), and Pacesetter[®]. The College Board is committed to the principles of equity and excellence, and that commitment is embodied in all of its programs, services, activities, and concerns.

Copyright © 2001 by College Entrance Examination Board. All rights reserved. College Board, Advanced Placement Program, AP, and the acorn logo are registered trademarks of the College Entrance Examination Board.

PHYSICS C

SECTION II, ELECTRICITY AND MAGNETISM

Time—45 minutes

3 Questions


Directions: Answer all three questions. The suggested time is about 15 minutes for answering each of the questions, which are worth 15 points each. The parts within a question may not have equal weight. Show all your work in this booklet in the spaces provided after each part, NOT in the green insert.



E&M 1. An isolated conducting sphere of radius $a = 0.20$ m is at a potential of $-2,000$ V.

- (a) Determine the charge Q_0 on the sphere.

$$-2000 = \frac{kQ_0}{a} \quad Q_0 = -4.44 \times 10^{-9} \text{ C}$$



The charged sphere is then concentrically surrounded by two uncharged conducting hemispheres of inner radius $b = 0.40$ m and outer radius $c = 0.50$ m, which are joined together as shown above, forming a spherical capacitor. A wire is connected from the outer sphere to ground, and then removed.

- (c) Determine the magnitude of the potential difference between the sphere and the conducting shell.

$$V_a - V_b = \int_a^b E_r dr = \frac{Q_0}{4\pi\epsilon_0} \int_a^b \frac{1}{r^2} dr = \left(\frac{1}{a} - \frac{1}{b}\right) \frac{Q_0}{4\pi\epsilon_0}$$
$$= \boxed{-999 \text{ V}}$$

- (d) Determine the capacitance of the spherical capacitor.

$$C = \frac{Q}{V} = \frac{4.44 \times 10^{-8}}{999} = 4.44 \times 10^{-11} \text{ F}$$

PHYSICS C

SECTION II, ELECTRICITY AND MAGNETISM

Time—45 minutes

3 Questions

Directions: Answer all three questions. The suggested time is about 15 minutes for answering each of the questions, which are worth 15 points each. The parts within a question may not have equal weight. Show all your work in this booklet in the spaces provided after each part, NOT in the green insert.




E&M 1. An isolated conducting sphere of radius $a = 0.20$ m is at a potential of $-2,000$ V.

(a) Determine the charge Q_0 on the sphere.

$$\left(\frac{1}{4\pi\epsilon_0}\right) \frac{q}{(0.20)} = -2000$$

$$\frac{1}{4\pi\epsilon_0} q = -400$$

$$q = -4.45 \times 10^{-8} \text{ C} = \boxed{-44.5 \text{ nC}}$$



The charged sphere is then concentrically surrounded by two uncharged conducting hemispheres of inner radius $b = 0.40$ m and outer radius $c = 0.50$ m, which are joined together as shown above, forming a spherical capacitor. A wire is connected from the outer sphere to ground, and then removed.

(b) Determine the magnitude of the electric field in the following regions as a function of the distance r from the center of the inner sphere.

i. $r < a$

INSIDE CONDUCTOR $\therefore E = 0$

ii. $a < r < b$

$\epsilon_0 \Phi = q$
 $\epsilon_0 \oint E \cdot dA = q$
 $\epsilon_0 (E) (4\pi r^2) = q$
 $E = \frac{q}{4\pi \epsilon_0 r^2}$
 $E = -\frac{dV}{dr} \therefore E = \frac{q}{4\pi \epsilon_0} \int_a^b \frac{dr}{r^2} = \frac{q}{4\pi \epsilon_0} \left[-\frac{1}{r} \right]_a^b = \frac{q}{4\pi \epsilon_0} \left[\frac{1}{b} - \frac{1}{a} \right]$

$E = \frac{1}{4\pi \epsilon_0} \left(\frac{-44.5 \times 10^{-9}}{.5} \right) = \boxed{-1000.14 \text{ N/C}}$

GO TO PART C

iii. $b < r < c$

INSIDE CONDUCTOR $\therefore E = 0$

iv. $r > c$

~~$E = \frac{q}{4\pi \epsilon_0 r^2} = \frac{q}{4\pi \epsilon_0} \int_a^b \frac{dr}{r^2} = \frac{q}{4\pi \epsilon_0} \left[-\frac{1}{r} \right]_a^b = \frac{q}{4\pi \epsilon_0} \left[\frac{1}{b} - \frac{1}{a} \right]$~~

$E = \frac{1}{4\pi \epsilon_0} \left(\frac{-44.5 \times 10^{-9}}{.5} \right) = \boxed{-800.11 \text{ N/C}}$

- (c) Determine the magnitude of the potential difference between the sphere and the conducting shell.

SEE b(ii)

$$\frac{q}{4\pi\epsilon_0} \left[\frac{1}{b} - \frac{1}{a} \right]$$

- (d) Determine the capacitance of the spherical capacitor.

$$C = \frac{q}{V}$$

$$= \frac{q}{\frac{q}{4\pi\epsilon_0} \left[\frac{1}{b} - \frac{1}{a} \right]} = \frac{4\pi\epsilon_0}{\left[\frac{1}{b} - \frac{1}{a} \right]}$$

PHYSICS C

SECTION II, ELECTRICITY AND MAGNETISM

Time—45 minutes

3 Questions

Directions: Answer all three questions. The suggested time is about 15 minutes for answering each of the questions, which are worth 15 points each. The parts within a question may not have equal weight. Show all your work in this booklet in the spaces provided after each part, NOT in the green insert.




E&M 1. An isolated conducting sphere of radius $a = 0.20$ m is at a potential of $-2,000$ V.

- (a) Determine the charge Q_0 on the sphere.

$$Ea = V = -2000 = 0.2E, E = -10,000 \text{ N/C}$$

$$-20,000 = \frac{Q_0}{4\pi\epsilon_0 a^2}, Q_0 = -4.448 \times 10^{-8} \text{ C}$$



The charged sphere is then concentrically surrounded by two uncharged conducting hemispheres of inner radius $b = 0.40$ m and outer radius $c = 0.50$ m, which are joined together as shown above, forming a spherical capacitor. A wire is connected from the outer sphere to ground, and then removed.

(b) Determine the magnitude of the electric field in the following regions as a function of the distance r from the center of the inner sphere.

i. $r < a$

$E = 0$, because of it being inside a conductor.

ii. $a < r < b$

$$E \cdot 4\pi r^2 = \frac{Q_0}{\epsilon_0} \quad E = \frac{Q_0}{4\pi\epsilon_0 r^2}$$

iii. $b < r < c$

$E = 0$, because of it being inside a conductor.

iv. $r > c$

$$E = \frac{Q_0}{4\pi\epsilon_0 r^2}$$

- (c) Determine the magnitude of the potential difference between the sphere and the conducting shell.

$$\Delta V = -\int_a^b E dr = -\int_a^b \frac{Q_0}{4\pi\epsilon_0 r^2} dr$$

$$\Delta V = -\frac{Q_0}{4\pi\epsilon_0} \int_a^b \frac{dr}{r^2}$$

$$\Delta V = \frac{Q_0}{4\pi\epsilon_0 r} \Big|_a^b$$

$$\Delta V = \frac{Q_0}{4\pi\epsilon_0} \left(\frac{1}{b} - \frac{1}{a} \right)$$

- (d) Determine the capacitance of the spherical capacitor.

$$Q_0 = CV, \quad C = \frac{Q_0}{V} = \frac{-4.448 \times 10^{-9} \text{ C}}{-2000 \text{ V}}$$

$$C = 2.224 \times 10^{-12} \text{ Farads}$$